

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number  
WO 01/00612 A2(51) International Patent Classification<sup>7</sup>: C07D 401/12,  
A61K 31/437, 31/4465, 31/4545, A61P 11/00, 31/12,  
C07D 471/04, 401/14 // (C07D 471/04, 235:00, 221:00)

(21) International Application Number: PCT/EP00/05675

(22) International Filing Date: 20 June 2000 (20.06.2000)

(25) Filing Language: English

(26) Publication Language: English



(30) Priority Data:  
99202088.3 28 June 1999 (28.06.1999) EP(71) Applicant (for all designated States except US):  
JANSSEN PHARMACEUTICA N.V. [BE/BE]; c/o  
De Corte, Filip - Ext. 3834, Patent Dept., Turnhoutseweg  
30, B-2340 Beerse (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JANSSENS, Frans,  
Eduard [BE/BE]; Janssen Pharmaceutica N.V., Turnhout-  
seweg 30, B-2340 Beerse (BE). MEERSMAN, Kathleen,Petrus, Marie-José [BE/BE]; Janssen Pharmaceutica  
N.V., Turnhoutseweg 30, B-2340 Beerse (BE). SOM-  
MEN, François, Maria [BE/BE]; Janssen Pharmaceutica  
N.V., Turnhoutseweg 30, B-2340 Beerse (BE). ANDRIES,  
Koenraad, Jozef, Lodewijk, Marcel [BE/BE]; Janssen  
Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse  
(BE).(74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica  
N.V., Patent Dept. - 3547, Turnhoutseweg 30, B-2340  
Beerse (BE).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,  
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,  
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European

[Continued on next page]

(54) Title: RESPIRATORY SYNCYTIAL VIRUS REPLICATION INHIBITORS



(57) Abstract: This invention concerns the compounds of formula (I), prodrugs, <I>N</I>-oxides, addition salts, quaternary amines, metal complexes or stereochemically isomeric forms thereof wherein -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- is a radical of formula -CH=CH-CH=CH-, -N=CH-CH=CH-, -CH=N-CH=CH-, -CH=CH-N=CH-, -CH=CH-CH=N- wherein each hydrogen atom may optionally be substituted; Q is a radical of formula (b-1), (b-2), (b-3), (b-4), (b-5), (b-6), (b-7), (b-8), wherein Alk is C<sub>1-6</sub>alkanediyl; Y<sup>1</sup> is a bivalent radical of formula -NR<sup>2</sup>- or -CH(NR<sup>2</sup>R<sup>4</sup>)-; X<sup>1</sup> is NR<sup>4</sup>, S, S(=O), S(=O)<sub>2</sub>, O, CH<sub>2</sub>, C(=O), CH(=CH<sub>2</sub>), CH(OH), CH(CH<sub>3</sub>), CH(OCH<sub>3</sub>), CH(SCH<sub>3</sub>), CH(NR<sup>5a</sup>R<sup>5b</sup>), CH<sub>2</sub>-NR<sup>4</sup> or NR<sup>4</sup>-CH<sub>2</sub>; X<sup>2</sup> is a direct bond, CH<sub>2</sub>, C(=O), NR<sup>4</sup>, C<sub>1-6</sub>alkyl-NR<sup>4</sup>, NR<sup>4</sup>-C<sub>1-6</sub>alkyl; t is 2 to 5; u is 1 to 5; v is 2 or 3; and whereby each hydrogen in Alk and in (b-3), (b-4), (b-5), (b-6), (b-7) and (b-8), may optionally be replaced by R<sup>3</sup>; provided that when R<sup>3</sup> is hydroxy or C<sub>1-6</sub>alkyloxy, then R<sup>3</sup> can not replace a hydrogen atom in the α position relative to a nitrogen atom; G is substituted C<sub>1-10</sub>alkanediyl wherein the substituent is attached via an oxygen atom; R<sup>1</sup> is an optionally substituted monocyclic heterocycle or aryl; R<sup>2</sup> is hydrogen, formyl, C<sub>1-6</sub>alkylcarbonyl, Hetcarbonyl, pyrrolidinyl, piperidinyl, homopiperidinyl, C<sub>3-7</sub>cycloalkyl or C<sub>1-10</sub>alkyl substituted with N(R<sup>6</sup>)<sub>2</sub> and optionally with another substituent; R<sup>3</sup> is hydrogen, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, arylC<sub>1-6</sub>alkyl or arylC<sub>1-6</sub>alkyloxy; R<sup>4</sup> is hydrogen, C<sub>1-6</sub>alkyl or arylC<sub>1-6</sub>alkyl; R<sup>5a</sup>, R<sup>5b</sup>, R<sup>5c</sup> and R<sup>5d</sup> are hydrogen or C<sub>1-6</sub>alkyl; or R<sup>5a</sup> and R<sup>5b</sup>, or R<sup>5c</sup> and R<sup>5d</sup> taken together form a bivalent radical of formula -(CH<sub>2</sub>)<sub>s</sub>- wherein s is 4 or 5; R<sup>6</sup> is hydrogen, C<sub>1-6</sub>alkyl, formyl, hydroxyC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonyl or C<sub>1-6</sub>alkyloxycarbonyl; aryl is optionally substituted phenyl; Het is pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl; as respiratory syncytial virus replication inhibitors; their preparation, compositions containing them and their use as a medicine.

WO 01/00612 A2



patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

*For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

**Published:**

- *Without international search report and to be republished upon receipt of that report.*

## RESPIRATORY SYNCYTIAL VIRUS REPLICATION INHIBITORS

---

5 The present invention is concerned with benzimidazoles and imidazopyridines having antiviral activity, in particular, they have an inhibitory activity on the replication of the respiratory syncytial virus. It further concerns their preparation and compositions comprising them, as well as their use as a medicine.

10 Human RSV or Respiratory Syncytial Virus is a large RNA virus, member of the family of Paramyxoviridae, subfamily pneumovirinae together with bovine RSV virus. Human RSV is responsible for a spectrum of respiratory tract diseases in people of all ages throughout the world. It is the major cause of lower respiratory tract illness during infancy and childhood. Over half of all infants encounter RSV in their first year of life, and almost all within their first two years. The infection in young children can cause  
15 lung damage that persists for years and may contribute to chronic lung disease in later life (chronic wheezing, asthma). Older children and adults often suffer from a (bad) common cold upon RSV infection. In old age, susceptibility again increases, and RSV has been implicated in a number of outbreaks of pneumonia in the aged resulting in significant mortality.

20 Infection with a virus from a given subgroup does not protect against a subsequent infection with an RSV isolate from the same subgroup in the following winter season. Re-infection with RSV is thus common, despite the existence of only two subtypes, A and B.


25 Today only three drugs have been approved for use against RSV infection. Ribavirin, a nucleoside analogue, provides an aerosol treatment for serious RSV infection in hospitalized children. The aerosol route of administration, the toxicity (risk of teratogenicity), the cost and the highly variable efficacy limit its use. The other two  
30 drugs, RespiGam<sup>®</sup> and palivizumab, polyclonal and monoclonal antibody immunostimulants, are intended to be used in a preventive way.

Other attempts to develop a safe and effective RSV vaccine have all met with failure thus far. Inactivated vaccines failed to protect against disease, and in fact in some cases  
35 enhanced disease during subsequent infection. Live attenuated vaccines have been tried with limited success. Clearly there is a need for an efficacious non-toxic and easy to administer drug against RSV replication.

-2-

EP-A-0,005,138 , EP-A-0,099,139 , EP-A-0,145,037 , EP-A-0,144,101 ,  
 EP-A-0,151,826 , EP-A-0,151,824 , EP-A-0,232,937 , EP-A-0,295,742 , EP 0,297,661 ,  
 EP-A-0,307,014 , WO 92 01697 describe benzimidazole and imidazopyridine  
 5 substituted piperidine and piperazine derivatives as antihistaminics, antiallergics or  
 serotonin antagonists.

The present invention concerns the compounds of formula (I)



their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and  
 10 stereochemically isomeric forms wherein

-a¹=a²-a³=a⁴- represents a bivalent radical of formula

-CH=CH-CH=CH- (a-1);


-N=CH-CH=CH- (a-2);

-CH=N-CH=CH- (a-3);

15 -CH=CH-N=CH- (a-4); or


-CH=CH-CH=N- (a-5);

wherein each hydrogen atom in the radicals (a-1), (a-2), (a-3), (a-4) and (a-5) may  
 optionally be replaced by halo, C<sub>1-6</sub>alkyl, nitro, amino, hydroxy, C<sub>1-6</sub>alkyl-  
 oxy, polyhaloC<sub>1-6</sub>alkyl, carboxyl, aminoC<sub>1-6</sub>alkyl, mono- or di(C<sub>1-4</sub>alkyl)-  
 20 aminoC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, hydroxyC<sub>1-6</sub>alkyl, or a radical of  
 formula




wherein =Z is =O, =CH-C(=O)-NR<sup>5a</sup>R<sup>5b</sup>, =CH<sub>2</sub>, =CH-C<sub>1-6</sub>alkyl, =N-OH or  
 =N-O-C<sub>1-6</sub>alkyl;


25 Q is a radical of formula




(b-1)



(b-2)




(b-3)



(b-4)

-3-



wherein Alk is C<sub>1-6</sub>alkanediyl;

Y<sup>1</sup> is a bivalent radical of formula -NR<sup>2</sup>- or -CH(NR<sup>2</sup>R<sup>4</sup>)-;

X<sup>1</sup> is NR<sup>4</sup>, S, S(=O), S(=O)<sub>2</sub>, O, CH<sub>2</sub>, C(=O), C(=CH<sub>2</sub>), CH(OH), CH(CH<sub>3</sub>),  
 5 CH(OCH<sub>3</sub>), CH(SCH<sub>3</sub>), CH(NR<sup>5a</sup>R<sup>5b</sup>), CH<sub>2</sub>-NR<sup>4</sup> or NR<sup>4</sup>-CH<sub>2</sub>;

X<sup>2</sup> is a direct bond, CH<sub>2</sub>, C(=O), NR<sup>4</sup>, C<sub>1-4</sub>alkyl-NR<sup>4</sup>, NR<sup>4</sup>-C<sub>1-4</sub>alkyl;

t is 2, 3, 4 or 5;

u is 1, 2, 3, 4 or 5;

v is 2 or 3; and

- 10 wherebyeach hydrogen atom in Alk and the carbocycles and the heterocycles defined in radicals (b-3), (b-4), (b-5), (b-6), (b-7) and (b-8) may optionally be replaced by R<sup>3</sup>; with the proviso that when R<sup>3</sup> is hydroxy or C<sub>1-6</sub>alkyloxy, then R<sup>3</sup> can not replace a hydrogen atom in the α position relative to a nitrogen atom;

G is C<sub>1-10</sub>alkanediyl substituted with one or more hydroxy, C<sub>1-6</sub>alkyloxy,

- 15 arylC<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, arylC<sub>1-6</sub>alkylthio, HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, C<sub>1-6</sub>alkyloxy-(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>- or arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-;

R<sup>1</sup> is a monocyclic heterocycle or aryl; said heterocycle being selected from piperidinyl, piperazinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl,

- 20 oxadiazolyl; and each heterocycle may optionally be substituted with 1 or where possible more, such as 2, 3 or 4, substituents selected from halo, hydroxy, amino, cyano, carboxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, arylC<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyl, mono-or di(C<sub>1-6</sub>alkyl)amino, mono-or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonylamino,  
 25 C<sub>1-6</sub>alkyl-SO<sub>2</sub>-NR<sup>5c</sup>-, aryl-SO<sub>2</sub>-NR<sup>5c</sup>-, C<sub>1-6</sub>alkyloxy carbonyl, -C(=O)-NR<sup>5c</sup>R<sup>5d</sup>, HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, halo(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, C<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>- and mono-or di(C<sub>1-6</sub>alkyl)amino(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-; each n independently is 1, 2, 3 or 4;

R<sup>2</sup> is hydrogen, formyl, C<sub>1-6</sub>alkylcarbonyl, Hetcarbonyl, pyrrolidinyl, piperidinyl,

- 30 homopiperidinyl, C<sub>3-7</sub>cycloalkyl substituted with N(R<sup>6</sup>)<sub>2</sub>, or C<sub>1-10</sub>alkyl substituted with N(R<sup>6</sup>)<sub>2</sub> and optionally with a second, third or fourth substituent selected from amino, hydroxy, C<sub>3-7</sub>cycloalkyl, C<sub>2-5</sub>alkanediyl, piperidinyl, mono-or di(C<sub>1-6</sub>alkyl)amino, C<sub>1-6</sub>alkyloxy carbonylamino, aryl and aryloxy;

$R^3$  is hydrogen, hydroxy,  $C_{1-6}$ alkyl,  $C_{1-6}$ alkyloxy, aryl $C_{1-6}$ alkyl or aryl $C_{1-6}$ alkyloxy;  
 $R^4$  is hydrogen,  $C_{1-6}$ alkyl or aryl $C_{1-6}$ alkyl;  
 $R^{5a}$ ,  $R^{5b}$ ,  $R^{5c}$  and  $R^{5d}$  each independently are hydrogen or  $C_{1-6}$ alkyl; or  
 $R^{5a}$  and  $R^{5b}$ , or  $R^{5c}$  and  $R^{5d}$  taken together form a bivalent radical of formula  $-(CH_2)_s-$   
5 wherein s is 4 or 5;  
 $R^6$  is hydrogen,  $C_{1-4}$ alkyl, formyl, hydroxy $C_{1-6}$ alkyl,  $C_{1-6}$ alkylcarbonyl or  
 $C_{1-6}$ alkyloxycarbonyl;  
aryl is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents  
selected from halo, hydroxy,  $C_{1-6}$ alkyl, hydroxy $C_{1-6}$ alkyl, polyhalo $C_{1-6}$ alkyl, and  
10  $C_{1-6}$ alkyloxy;  
Het is pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl.

The term prodrug as used throughout this text means the pharmacologically acceptable  
derivatives, e.g. esters and amides, such that the resulting biotransformation product of  
15 the derivative is the active drug as defined in the compounds of formula (I). The  
reference by Goodman and Gilman (The Pharmacological Basis of Therapeutics, 8<sup>th</sup>  
ed., McGraw-Hill, Int. Ed. 1992, "Biotransformation of Drugs", p. 13-15) describing  
prodrugs generally, is hereby incorporated.

20 As used herein  $C_{1-3}$ alkyl as a group or part of a group defines straight or branched chain  
saturated hydrocarbon radicals having from 1 to 3 carbon atoms such as methyl, ethyl,  
propyl, 1-methylethyl and the like;  $C_{1-4}$ alkyl as a group or part of a group defines  
straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon  
atoms such as the group defined for  $C_{1-3}$ alkyl and butyl and the like;  $C_{2-4}$ alkyl as a  
25 group or part of a group defines straight or branched chain saturated hydrocarbon  
radicals having from 2 to 4 carbon atoms such as ethyl, propyl, 1-methylethyl, butyl and  
the like;  $C_{1-6}$ alkyl as a group or part of a group defines straight or branched chain  
saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the groups  
defined for  $C_{1-4}$ alkyl and pentyl, hexyl, 2-methylbutyl and the like;  $C_{1-9}$ alkyl as a group  
30 or part of a group defines straight or branched chain saturated hydrocarbon radicals  
having from 1 to 9 carbon atoms such as the groups defined for  $C_{1-6}$ alkyl and heptyl,  
octyl, nonyl, 2-methylhexyl, 2-methylheptyl and the like;  $C_{1-10}$ alkyl as a group or part of  
a group defines straight or branched chain saturated hydrocarbon radicals having from 1  
to 10 carbon atoms such as the groups defined for  $C_{1-9}$ alkyl and decyl, 2-methylnonyl  
35 and the like.  $C_{3-7}$ cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl,  
cyclohexyl and cycloheptyl;  $C_{2-5}$ alkanediyl defines bivalent straight and branched chain  
saturated hydrocarbon radicals having from 2 to 5 carbon atoms such as, for example,

1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl, 1,2-propanediyl, 2,3-butanediyl, 1,5-pentanediyl and the like, C<sub>2-5</sub>alkanediyl is substituted on C<sub>1-10</sub>alkyl as provided for in the definition of R<sup>2</sup>, it is meant to be substituted on one carbon atom thus forming a spiro moiety; C<sub>1-4</sub>alkanediyl defines bivalent straight and branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as, for example, methylene, 1,2-ethanediyl, 1,3-propanediyl, 1,4-butanediyl and the like; C<sub>1-6</sub>alkanediyl is meant to include C<sub>1-4</sub>alkanediyl and the higher homologues thereof having from 5 to 6 carbon atoms such as, for example, 1,5-pentanediyl, 1,6-hexanediyl and the like; C<sub>1-10</sub>alkanediyl is meant to include C<sub>1-6</sub>alkanediyl and the higher homologues thereof having from 7 to 10 carbon atoms such as, for example, 1,7-heptanediyl, 1,8-octanediyl, 1,9-nonanediyl, 1,10-decanediyl and the like.

As used herein before, the term (=O) forms a carbonyl moiety when attached to a carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety when two of said terms are attached to a sulfur atom. The term (=N-OH) forms a hydroxylimine moiety when attached to a carbon atom.

The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and hereinafter, polyhaloC<sub>1-6</sub>alkyl as a group or part of a group is defined as mono- or polyhalosubstituted C<sub>1-6</sub>alkyl, in particular methyl with one or more fluoro atoms, for example, difluoromethyl or trifluoromethyl. In case more than one halogen atoms are attached to an alkyl group within the definition of polyhaloC<sub>1-4</sub>alkyl, they may be the same or different.

When any variable (e.g. aryl, R<sup>2</sup>, R<sup>3</sup>, R<sup>4</sup>, R<sup>5a</sup>, R<sup>5b</sup> etc.) occurs more than one time in any constituent, each definition is independent.

It will be appreciated that some of the compounds of formula (I) and their prodrugs, N-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms which the compounds of formula (I), and their prodrugs, N-oxides, addition salts, quaternary amines, metal complexes or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers

of the basic molecular structure as well as each of the individual isomeric forms of formula (I) and their prodrugs, *N*-oxides, salts, solvates or quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers.

- 5 Stereochemically isomeric forms of the compounds of formula (I) are obviously intended to be embraced within the scope of this invention. As used hereinafter the terms R or S are well-known by the person skilled in the art

- For some of the compounds of formula (I), their prodrugs, *N*-oxides, salts, solvates, quaternary amines, or metal complexes and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined. In these cases the stereoisomeric form which was first isolated is designated as "A" and the second as "B", without further reference to the actual stereochemical configuration. However, said "A" and "B" stereoisomeric forms can be unambiguously characterized by for instance their optical rotation in case "A" and "B" have an enantiomeric relationship. A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction. In case "A" and "B" are stereoisomeric mixtures, they can be further separated whereby the respective first fractions isolated are designated "A1" and "B1" and the second as "A2" and "B2", without further reference to the actual stereochemical configuration.
- 10  
15  
20

- For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.
- 25

- The pharmaceutically acceptable acid and base addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid and base addition salt forms which the compounds of formula (I) are able to form. The pharmaceutically acceptable acid addition salts can conveniently be obtained by treating the base form with such appropriate acid. Appropriate acids comprise, for example, inorganic acids such as hydrohalic acids, *e.g.* hydrochloric or hydrobromic acid, sulfuric, nitric, phosphoric and the like acids; or organic acids such as, for example, acetic, propanoic, hydroxyacetic, lactic, pyruvic, oxalic (*i.e.* ethanedioic), malonic, succinic (*i.e.* butanedioic acid), maleic, fumaric, malic (*i.e.* hydroxybutanedioic acid), tartaric, citric,
- 30  
35



methanesulfonic, ethanesulfonic, benzenesulfonic, *p*-toluenesulfonic, cyclamic, salicylic, *p*-aminosalicylic, pamoic and the like acids.

Conversely said salt forms can be converted by treatment with an appropriate base into the free base form.

The compounds of formula (I) containing an acidic proton may also be converted into their non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. the benzathine, *N*-methyl-D-glucamine, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like.

The term addition salt as used hereinabove also comprises the solvates which the compounds of formula (I) as well as the salts thereof, are able to form. Such solvates are for example hydrates, alcoholates and the like.

The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl *p*-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.

It will be appreciated that the compounds of formula (I) may have metal binding, chelating, complexing properties and therefore may exist as metal complexes or metal chelates. Such metalated derivatives of the compounds of formula (I) are intended to be included within the scope of the present invention.

Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included within the scope of the present invention.

-8-

A special group of compounds are those compounds of formula (I) wherein one or more of the following restrictions apply:

- Q is a radical of formula (b-1), (b-3), (b-4), (b-5), (b-6), (b-7) or (b-8);
- X<sup>2</sup> is a direct bond, CH<sub>2</sub> or C(=O);
- 5 - R<sup>2</sup> is hydrogen, pyrrolidinyl, piperidinyl, homopiperidinyl, C<sub>3-7</sub>cycloalkyl substituted with NHR<sup>6</sup>, or C<sub>1-10</sub>alkyl substituted with NHR<sup>6</sup> and optionally with a second, third or fourth substituent selected from amino, hydroxy, C<sub>3-7</sub>cycloalkyl, C<sub>2-5</sub>alkanediyl, piperidinyl, mono- or di(C<sub>1-6</sub>alkyl)amino, C<sub>1-6</sub>alkyloxycarbonylamino, aryl and aryloxy;
- R<sup>3</sup> is hydrogen, hydroxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy or arylC<sub>1-6</sub>alkyl;
- 10 - R<sup>6</sup> is hydrogen, C<sub>1-4</sub>alkyl, formyl, C<sub>1-6</sub>alkylcarbonyl or C<sub>1-6</sub>alkyloxycarbonyl.

Also an interesting group of compounds are those compounds of formula (I) wherein one or more of the following restrictions apply :

- a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- is a radical of formula (a-1) or (a-2);
- 15 R<sup>1</sup> is phenyl optionally substituted with halo, C<sub>1-6</sub>alkyl or C<sub>1-4</sub>alkyloxy; or pyridyl optionally substituted with 1 or more substituents selected from arylC<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, aryl, mono- or di(C<sub>1-6</sub>alkyl)amino, C(=O)-NR<sup>5c</sup>R<sup>5d</sup>, halo or C<sub>1-6</sub>alkyl;
- G is C<sub>1-4</sub>alkanediyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-,
- 20 C<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>- or arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-;
- Q is a radical of formula (b-5) wherein v is 2, and Y<sup>1</sup> is N-R<sup>2</sup>;
- X<sup>1</sup> is NH or CH<sub>2</sub>;
- R<sup>2</sup> is hydrogen or C<sub>1-10</sub>alkyl substituted with NHR<sup>6</sup> wherein R<sup>6</sup> is hydrogen or C<sub>1-6</sub>alkyloxycarbonyl.

25

Particular compounds are those compounds of formula (I) wherein R<sup>2</sup> is C<sub>1-10</sub>alkyl substituted with NH<sub>2</sub>.

- Other particular compounds are those compounds of formula (I) wherein G is
- 30 methylene or 1,2-ethanediyl, both substituted with hydroxy, C<sub>1-6</sub>alkyloxy, HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, C<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>- or arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-.

- Also particular compounds are those compounds of formula (I) wherein R<sup>1</sup> is pyridyl, preferably 2-pyridyl, substituted with one or 2 substituents selected from halo, hydroxy,
- 35 amino, cyano, carboxy, C<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkylthio, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, aryl, arylC<sub>1-6</sub>alkyl, arylC<sub>1-6</sub>alkyloxy, hydroxyC<sub>1-6</sub>alkyl, mono- or di(C<sub>1-6</sub>alkyl)amino, mono- or di(C<sub>1-6</sub>alkyl)aminoC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkylcarbonylamino, C<sub>1-6</sub>alkyl-SO<sub>2</sub>-NR<sup>5c</sup>-, aryl-SO<sub>2</sub>-NR<sup>5c</sup>-, C<sub>1-6</sub>alkyloxycarbonyl, -C(=O)-NR<sup>5c</sup>R<sup>5d</sup>,

HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, halo(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, C<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>- and mono-or di(C<sub>1-6</sub>alkyl)amino(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, preferably selected from arylC<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, aryl, mono-or di(C<sub>1-6</sub>alkyl)amino, C(=O)-NR<sup>5a</sup>R<sup>5b</sup>, halo or C<sub>1-6</sub>alkyl.

5

Preferred compounds are those compounds of formula (I) wherein R<sup>1</sup> is an optionally substituted 2-pyridyl moiety, in particular, a 2-pyridyl, a 6-substituted-2-pyridyl or a 3,6-disubstituted-2-pyridyl moiety.

10

Preferred compounds are

[(A),(S)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1H-benzimidazol-2-amine (compound 69);

[(A),(S)]-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 75);

15

(±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 86);

N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-6-chloro-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-4-methyl-1H-benzimidazol-2-amine trihydrochloride trihydrate (compound 88);

20

[(A),(R)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate (compound 68);

(±)-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 12);

25

[(A),(S)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate (compound 67);

(±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 83);

[(A),(R)]-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate (compound 74);

30

(±)-N-[1-(2-aminoethyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-2-benzimidazol-2-amine (compound 9);

(±)-N-[1-(2-aminoethyl)-4-piperidinyl]-1-[(2-ethoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 64);

[(B),(S)] N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate (compound 76);

35

(±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-3-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-7-methyl-3H-imidazo[4,5-b]pyridin-2-amine (compound 89);

(±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(2-ethoxyethoxy)(6-phenyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine (compound 85);

(±)-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine (compound 82);

- 5 the prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms thereof.

Most preferred are

(±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-4-methyl-1*H*-benzimidazol-2-amine monohydrate (compound 87);

10 [(A),(R))-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-amine (compound 70);


(±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-amine (compound 10);

- 15 the prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms thereof.

In general, compounds of formula (I) can be prepared by reacting an intermediate of formula (II-a) or (II-b), wherein P represents a protecting group, such as, for example


20 C<sub>1-4</sub>alkyloxycarbonyl, or those protecting groups mentioned in Chapter 7 of 'Protective Groups in Organic Synthesis' by T Greene and P. Wuyts (John Wiley & Sons Inc., 1991), with an intermediate of formula (III), wherein W<sub>1</sub> is a suitable leaving group, such as a halo atom, e.g. chloro, bromo, in the presence of a suitable base, such as, e.g. sodium hydride. Said reaction can be performed in a reaction-inert solvent, such as

25 *N,N*-dimethylformamide.




Compounds of formula (I) wherein, in the definition of Q, R<sup>2</sup> or at least one R<sup>6</sup> substituent is hydrogen, said Q being represented by H-Q<sub>1</sub>, and said compounds being represented by formula (I-a), can be prepared by deprotecting an intermediate of formula (IV) wherein P represents a protecting group, for example C<sub>1</sub>-

- 5 4alkyloxycarbonyl, benzyl, or those protecting groups mentioned in Chapter 7 of 'Protective Groups in Organic Synthesis' by T Greene and P. Wuyts (John Wiley & Sons Inc., 1991).




- When P represents, for example, C<sub>1-4</sub>alkyloxycarbonyl, said deprotection reaction can be performed by, for example, acidic hydrolysis in the presence of a suitable acid, such as hydrobromic, hydrochloric, sulfuric, acetic, or trifluoroacetic acid or a mixture of said acids, or by alkaline hydrolysis in the presence of a suitable base, such as, for example potassium hydroxide, in a suitable solvent such as water, alcohol, a mixture of water-alcohol, methylene chloride. Suitable alcohols are methanol, ethanol,
- 15 2-propanol, 1-butanol and the like. In order to enhance the rate of the reaction, it is advantageous to heat the reaction mixture, in particular up to the reflux temperature. Alternatively, when P represents, for example, benzyl, the deprotection reaction can be performed by catalytic hydrogenation in the presence of hydrogen and an appropriate catalyst in a reaction-inert solvent. A suitable catalyst in the above reaction is, for example, platinum-on-charcoal, palladium-on-charcoal, and the like. An appropriate
- 20 reaction-inert solvent for said reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like, an ester, e.g. ethylacetate and the like, an acid, e.g. acetic acid and the like.
- 25 The catalytic hydrogenation reaction described above can also be used to prepare a compound of formula (I-a) by deprotecting and reducing an intermediate of formula (IV) wherein Q<sub>1</sub> comprises an unsaturated bond, said Q<sub>1</sub> being represented by Q<sub>1a</sub>(CH=CH), and said intermediate being represented by formula (IV-a).

-12-



Compounds of formula (I) wherein, in the definition of Q, both R<sup>6</sup> substituents are hydrogen or R<sup>2</sup> and R<sup>4</sup> are both hydrogen, said Q being represented by H<sub>2</sub>N-Q<sub>2</sub>, and said compounds being represented by formula (I-a-1), can also be prepared by


5 deprotecting an intermediate of formula (V).



Said deprotection reaction can be performed in the presence of a suitable base such as, for example hydrazine, or in the presence of a suitable acid, such as hydrochloric acid and the like, in a suitable solvent, such as an alcohol, acetic acid and the like.


10

Compounds of formula (I-a-1) can also be prepared by deprotecting an intermediate of formula (VI) according to the procedure described for the preparation of compounds of formula (I-a).




15 Compounds of formula (I-a) or (I-a-1), wherein Q<sub>1</sub> or Q<sub>2</sub> comprise a hydroxy substituent, said Q<sub>1</sub> or Q<sub>2</sub> being represented by Q<sub>1</sub>·(OH) or Q<sub>2</sub>·(OH), and said compounds being represented by formula (I-a-2) or (I-a-1-1), can be prepared by deprotecting an intermediate of formula (VII) or (VIII) as described hereinabove for the preparation of compounds of formula (I-a).

-13-




- Compounds of formula (I) wherein, in the definition of Q, both  $R^6$  substituents are hydrogen or  $R^2$  and  $R^4$  are both hydrogen, and the carbon adjacent to the nitrogen carrying the  $R^6$ , or  $R^2$  and  $R^4$  substituents contains at least one hydrogen, said Q being represented by  $H_2N-Q_3H$ , and said compounds being represented by formula (I-a-1-2) can also be obtained by reductive amination of intermediates of formula (IX) in the presence of a suitable amination reagent, such as, for example, ammonia, hydroxylamine, or benzylamine, and in the presence of a suitable reducing agent, e.g. hydrogen, and an appropriate catalyst. An appropriate catalyst in the above reaction is, for example, platinum-on-charcoal, palladium-on-charcoal, rhodium-on- $Al_2O_3$ , and the like, optionally in the presence of a catalyst poison, such as a thiophene solution. A suitable reaction-inert solvent for the above reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like.




Compounds of formula (I), wherein Q comprises a  $-CH_2NH_2$  moiety, said Q being represented by  $H_2N-CH_2-Q_4$ , and said compounds being represented by formula (I-a-1-3) can be prepared by reducing an intermediate of formula (X).


-14-



- Said reduction can be performed with a suitable reducing agent, such as lithium aluminium hydride or hydrogen, optionally in the presence of a suitable catalyst, such as Raney Nickel. A suitable solvent for the above reaction is, for example,
- 5 tetrahydrofuran, or a solution of ammonia in an alcohol. Suitable alcohols are methanol, ethanol, 2-propanol and the like. Said reduction reaction performed in a solution of ammonia in an alcohol can also be used to prepare compounds of formula (I-a-1-3), wherein  $R^1$  is substituted with  $C_{1-6}$ alkyloxy $C_{1-6}$ alkyl, said  $R^1$  being represented by  $R^1-C_{1-6}$ alkyloxy $C_{1-6}$ alkyl, and said compounds being represented by
- 10 formula (I-a-1-3-1) starting from an intermediate of formula (X-a).




- Compounds of formula (I), wherein Q comprises a  $-CH_2-CHOH-CH_2-NH_2$  moiety, said Q being represented by  $H_2N-CH_2-CHOH-CH_2-Q_4$ , and said compounds being represented by formula (I-a-1-3-2), can be prepared by reacting an intermediate of
- 15 formula (XI) with ammonia in the presence of a suitable reaction-inert solvent, such as an alcohol, e.g. methanol.




- Compounds of formula (I), wherein, in the definition of Q,  $R^2$  or one  $R^6$  substituent is formyl, said Q being represented by  $H-C(=O)-Q_1$ , and said compounds being
- 20 represented by formula (I-b), can be prepared by reacting an intermediate of formula (XII) with formic acid, formamide and ammonia.




-15-



- Compounds of formula (I), wherein, in the definition of Q, R<sup>2</sup> is other than hydrogen, said R<sup>2</sup> being represented by R<sup>2a</sup>, R<sup>4</sup> is hydrogen, and the carbon atom adjacent to the nitrogen atom carrying the R<sup>2</sup> and R<sup>4</sup> substituents, carries also at least one hydrogen atom, said Q being represented by R<sup>2a</sup>-NH-HQ<sub>5</sub>, and said compounds being represented by formula (I-c), can be prepared by reductive amination of an intermediate of formula (XIII) with an intermediate of formula (XIV) in the presence of a suitable reducing agent, such as hydrogen, and a suitable catalyst, such as palladium-on-charcoal, platinum-on-charcoal, and the like. A suitable reaction-inert solvent for the above reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like.



- Compounds of formula (I-c), wherein R<sup>2a</sup> represents C<sub>1-10</sub>alkyl substituted with N(R<sup>6</sup>)<sub>2</sub> and with hydroxy, and the carbon atom carrying the hydroxy, carries also two hydrogen atoms, said R<sup>2a</sup> being represented by [(C<sub>1-9</sub>alkyl)CH<sub>2</sub>OH]-N(R<sup>6</sup>)<sub>2</sub>, and said compounds being represented by formula (I-c-1), can be prepared by reducing an intermediate of formula (XV) in the presence of a suitable reducing agent, such as lithium aluminium hydride, in a suitable reaction-inert solvent, such as tetrahydrofuran.






- Compounds of formula (I) wherein, in the definition of Q, R<sup>2</sup> or one R<sup>6</sup> substituent is hydrogen, said Q being represented by H-Q<sub>1</sub>, and wherein R<sup>1</sup> is aryl or a monocyclic heterocycle substituted with 1 or more substituents selected from hydroxy, hydroxyC<sub>1-6</sub>alkyl, or HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, said substituents being represented by formula A-OH, said R<sup>1</sup> being represented by R<sup>1a</sup>-(A-OH)<sub>w</sub>, with w being the amount of

-16-


substituents on  $R^{1a}$  ranging from 1 to 4, and said compounds being represented by formula (I-d), can be prepared by deprotecting an intermediate of formula (XVI) with a suitable acid, such as hydrochloric acid and the like, optionally in the presence of a suitable solvent, such as an alcohol. Suitable alcohols are methanol, ethanol, 2-propanol and the like.

Alternatively, one protecting group may also protect more than one substituent of  $R^{1a}$ , said protecting group being represented by  $P_1$ , as represented by formula (XVI-a). The two ways of protecting the substituents of  $R^{1a}$ , i.e. with a separate, as in formula (XVI), or a combined, as in formula (XVI-a), protecting group, may also be combined in the same intermediate, as represented by formula (XVI-b).




Compounds of formula (I), wherein Q is a radical of formula (b-2), said compounds being represented by formula (I-e), can be prepared by reacting an intermediate of formula (XVII) with an intermediate of formula (XVIII) in the presence of sodium cyanide and a suitable reaction-inert solvent, such as an alcohol, e.g. methanol and the like.


-17-



- Compounds of formula (I), wherein in the definition of Q, X<sup>2</sup> is C<sub>2-4</sub>alkyl-NR<sup>4</sup>, said Q being represented by Q<sub>6</sub>N-CH<sub>2</sub>-C<sub>1-3</sub>alkyl-NR<sup>4</sup>, and said compounds being represented by formula (I-p), can be prepared by reacting an intermediate of formula (XIX) with an intermediate of formula (XX) in the presence of isopropyl titanate (IV) and a suitable reducing agent, such as NaBH<sub>3</sub>CN, and in the presence of a suitable reaction-inert solvent, such as methylene chloride and an alcohol, e.g. ethanol.




- Compounds of formula (I-p), wherein R<sup>2</sup> is C<sub>1-6</sub>alkylcarbonyl, and Q is a radical of formula (b-6), wherein Y<sup>1</sup> is NR<sup>2</sup>, said compounds being represented by formula (I-p-1), can be prepared by reacting an intermediate of formula (XIX) with an intermediate of formula (XX-a) according to the procedure described for the preparation of a compound of formula (I-p).




- Compounds of formula (I), wherein G is substituted with hydroxy or HO(-CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-, said G being represented by G<sub>1</sub>-OH, and said compounds being represented by formula (I-q), may be prepared by deprotecting an intermediate of formula (XXI), wherein P represents a suitable protecting group, for example, benzyl. Said deprotection reaction

can be performed by catalytic hydrogenation in the presence of hydrogen and an appropriate catalyst in a reaction-inert solvent. A suitable catalyst in the above reaction is, for example, platinum-on-charcoal, palladium-on-charcoal, and the like. An appropriate reaction-inert solvent for said reaction is, for example, an alcohol, e.g. methanol, ethanol, 2-propanol and the like, an ester, e.g. ethylacetate and the like, an acid, e.g. acetic acid and the like.



Compounds of formula (I), wherein G is substituted with hydroxy and the carbon atom carrying the hydroxy substituent carries also at least one hydrogen, said G being represented by H-G<sub>2</sub>-OH, and said compounds being represented by formula (I-q-1), can also be prepared by reducing an intermediate of formula (XXII).




Said reduction reaction can be performed in the presence of a suitable reducing agent, such as, for example sodium borohydride, in a reaction-inert solvent, such as an alcohol or tetrahydrofuran or a mixture thereof. Suitable alcohols are methanol, ethanol, 2-propanol and the like.

Compounds of formula (I) may be converted into each other following art-known functional group transformation reactions, comprising those described hereinafter.


The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboxy-

oxoic acid or halo substituted benzenecarboxylic acid, e.g. 3-chlorobenzenecarboxylic acid, peroxoalkanoic acids, e.g. peroxoacetic acid, alkylhydroperoxides, e.g. t.butyl hydro-peroxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated  
 5 hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

Compounds of formula (I), wherein  $R^1$  is monocyclic heterocycle substituted with  $C_{1-6}$ alkyloxycarbonyl, said  $R^1$  being represented by  $R^{1'}-C(=O)OC_{1-6}alkyl$ , and said compounds being represented by formula (I-f), can be prepared by esterification of a  
 10 compound of formula (I-g) in the presence of a suitable alcohol, e.g. methanol, ethanol, propanol, butanol, pentanol, hexanol and the like, and in the presence of a suitable acid, such as hydrochloric acid and the like.




Compounds of formula (I-a) may be converted into compounds of formula (I) wherein, in the definition of Q,  $R^2$  or at least one  $R^6$  substituent is other than hydrogen, said  $R^2$  or  $R^6$  being represented by  $Z_1$ , said Q being represented by  $Z_1-Q_1$ , and said compounds being represented by formula (I-h), by reaction with a reagent of formula (XXIII), wherein  $W_2$  is a suitable leaving group, such as a halo atom, e.g. bromo, or 4-methylbenzenesulphonate, in the presence of a suitable base, such as, for example  
 20 disodium carbonate, dipotassium carbonate, sodium hydroxide and the like, in a reaction-inert solvent, e.g. 3-methyl-2-butanone, acetonitrile, *N,N*-dimethylformamide.




Compounds of formula (I-h), wherein, in the definition of  $Z_1$ ,  $R^2$  is  $CH_2-C_{1-9}alkyl$  substituted with  $N(R^6)_2$ , said compounds being represented by formula (I-h-1), can also  
 25 be prepared by reacting a compound of formula (I-a) wherein, in the definition of H-Q<sub>1</sub>,  $R^2$  is hydrogen, said H-Q<sub>1</sub> being represented by H-Q<sub>1b</sub>, and said compounds being represented by formula (I-a-3), with an intermediate of formula (XXIV), in the presence


of a suitable reducing agent, such as sodium cyanoborohydride, in a suitable reaction-inert solvent, such as an alcohol.



- Compounds of formula (I-h), wherein  $Z_1$  comprises formyl,  $\text{C}_{1-6}$ alkylcarbonyl, Hetcarbonyl or  $\text{C}_{1-6}$ alkyloxycarbonyl, said  $Z_1$  being represented by  $Z_{1a}$ , and said compounds being represented by formula (I-h-2), can be converted into compounds of formula (I-a), by acidic hydrolysis in the presence of a suitable acid, such as hydrobromic, hydrochloric, sulfuric, acetic, or trifluoroacetic acid or a mixture of said acids, or by alkaline hydrolysis in the presence of a suitable base, such as, for example potassium hydroxide, in a suitable solvent such as water, alcohol, a mixture of water-alcohol, methylene chloride. Suitable alcohols are methanol, ethanol, 2-propanol, 1-butanol, sec. butanol and the like. In order to enhance the rate of the reaction, it is advantageous to work at elevated temperatures.




- Compounds of formula (I-b) can be prepared by reacting a compound of formula (I-a) with formic acid.




- Compounds of formula (I) wherein  $\text{R}^1$  is monocyclic heterocycle or aryl substituted with hydroxy, said  $\text{R}^1$  being represented by  $\text{HO}-\text{R}^{1'}$ , and said compounds being represented by formula (I-i), can be prepared by deprotecting a compound of formula (I-j), wherein  $\text{R}^1$  is monocyclic heterocycle or aryl substituted with  $\text{C}_{1-6}$ alkyloxy or aryl $\text{C}_{1-6}$ alkyloxy, said  $\text{C}_{1-6}$ alkyl or aryl $\text{C}_{1-6}$ alkyl being represented by  $Z_2$ , and said  $\text{R}^1$

-21-

being represented by  $Z_2-O-R^1$ . Said deprotection can be performed in a reaction-inert solvent, such as, for example methylene chloride, in the presence of a suitable deprotecting agent, e.g. tribromoborane.



- 5 Compounds of formula (I) wherein  $R^1$  is monocyclic heterocycle substituted with halo( $-\text{CH}_2-\text{CH}_2-\text{O}$ ) $_n$ , said compounds being represented by formula (I-k), can be converted into compounds of formula (I-l-1) or (I-l-2) by reaction with an appropriate amine of formula (XXV) or (XXVI) in a suitable reaction-inert solvent, e.g. tetrahydrofuran.




10


Compounds of formula (I) wherein  $R^1$  is monocyclic heterocycle or aryl substituted with halo, said compounds being represented by formula (I-m) can be converted into compounds of formula (I) by reaction with 1-butanethiol in the presence of palladium-on-charcoal and CaO in a suitable reaction-inert solvent, such as

15 tetrahydrofuran.

-22-



- Compounds of formula (I) wherein a hydrogen atom in the radicals of formula (a-1), (a-2), (a-3), (a-4) or (a-5) is replaced by nitro, said compounds being represented by formula (I-n) may be reduced to a compound of formula (I-o) in the presence of a suitable reducing agent, such as hydrogen, optionally in the presence of a suitable catalyst, such as platinum-on-charcoal, and optionally in the presence of a suitable catalyst poison, e.g. a thiophene solution. The reaction may be performed in a suitable reaction-inert solvent, such as an alcohol.




- In the following paragraphs, there are described several methods of preparing the intermediates in the foregoing preparations. A number of intermediates and starting materials are commercially available or are known compounds which may be prepared according to conventional reaction procedures generally known in the art or analogous to the procedures described in EP-A-0005318, EP-A-0099139, EP-A-0151824 , EP-A-0151826 , EP-A-0232937 , EP-A-0295742 , EP-A-0297661 , EP-A-0539420 , EP-A-0539421 , US 4,634,704 , US 4,695,569.

- In the foregoing and the following preparations, the reaction mixture is worked up following art-known methods and the reaction product is isolated and, if necessary, further purified.


- Intermediates of formula (III) can be prepared by reacting an intermediate of formula (XXVII) with a suitable leaving group, i.e. W<sub>1</sub>, introducing agent, e.g. 1-halo-2,5-pyrrolidinedione in the presence of dibenzoyl peroxide, in a reaction-inert solvent, e.g. tetrachloromethane.




-23-




- Intermediates of formula (XXVII), wherein  $\text{R}^1$  is monocyclic heterocycle or aryl substituted with chloro, said  $\text{R}^1$  being represented by  $\text{Cl-R}^1$  and said intermediates being represented by formula (XXVII-a) can be prepared by reacting an intermediate of formula (XXVIII), wherein  $(\text{O=})\text{R}^{1b}\text{H}$  is defined as a carbonyl derivative of  $\text{R}^1$  wherein one carbon or nitrogen, adjacent to the carbonyl, carries at least one hydrogen, with phosphorus oxychloride. Intermediates of formula (XXVIII) may also react as their enol tautomeric forms.



- Intermediates of formula (III) wherein  $\text{W}_1$  is chloro, which is attached to a carbon atom carrying at least one hydrogen, said  $\text{G}$  being represented by  $\text{G}_3\text{H}$ , and said intermediates being represented by formula (III-a) can also be prepared by reacting an intermediate of formula (XXIX) with thionylchloride in a reaction-inert solvent, e.g. methylenechloride.



- Intermediates of formula (XXIX) can be prepared by reducing an intermediate of formula (XXX) in a reaction-inert solvent, e.g. an alcohol, in the presence of a suitable reducing agent, e.g. sodium borohydride.



- Alternatively, intermediates of formula (XXIX) can also be prepared by deprotecting an intermediate of formula (XXXI), wherein  $\text{P}$  is a suitable protecting group, e.g.  $\text{C}_{1-4}$ alkylcarbonyl, in a reaction-inert solvent, such as an alcohol, in the presence of a suitable base, e.g. sodium hydroxide.

-24-



(XXXI)

(XXIX)

Intermediates of formula (XXX), wherein  $G_3(=O)$  is  $CH(=O)$ , said intermediates being represented by formula (XXX-a), can be prepared by reacting an intermediate of formula (XXXII), wherein  $W_3$  is a suitable leaving group, such as a halo atom, e.g.


- 5 bromo, with *N,N*-dimethylformamide in the presence of butyllithium in a reaction-inert solvent, e.g. tetrahydrofuran, diethylether or a mixture thereof.




(XXXII)

(XXX-a)


- Intermediates of formula (IV) can be prepared by reacting an intermediate of formula (XXXIII-a) or (XXXIII-b), wherein P represents a suitable protecting group, such as, for example,  $C_{1-4}$ alkyloxycarbonyl, with an intermediate of formula (III) according to the reaction described for the general preparation of compounds of formula (I).




- Intermediates of formula (IV) can also be prepared by reacting an intermediate of formula (XXXIII-a) with an intermediate of formula (XXXIV) that has reacted with methanesulfonyl chloride, in the presence of a suitable base, such as sodium hydride, and in the presence of a suitable reaction-inert solvent, e.g. *N,N*-dimethylformamide.




Intermediates of formula (IV) can also be prepared by a cyclization reaction of an intermediate of formula (XXXV) in a reaction-inert solvent, e.g. an alcohol or *N,N*-dimethylformamide, in the presence of mercury oxide and sulphur.



- 5 Intermediates of formula (IV) wherein  $Q_1$  comprises an unsaturated bond, said  $Q_1$  being represented by  $Q_{1a}(CH=CH)$ , and said intermediates by formula (IV-a), can be prepared by reacting an intermediate of formula (XXXVI) with an intermediate of formula (III) in the presence of a suitable base, such as dipotassium carbonate.




- 10 Intermediates of formula (IV) wherein, in the definition of  $Q_1$ , the  $X^1$  or  $X^2$  moieties in the radicals of formula (b-1) to (b-8) represent NH, said  $Q_1$  being represented by  $Q_{1c}-NH$ , and said intermediates by formula (IV-b), may also be prepared by reacting an intermediate of formula (XXXVII) with an intermediate of formula (XXXVIII).




- 15 Intermediates of formula (IV) wherein  $R^1$  is monocyclic heterocycle substituted with amino or mono- or di( $C_{1-6}$ alkyl)amino, said  $R^1$  being represented by  $R^{5a}R^{5b}N-R^1$ , wherein  $R^{5a}$  and  $R^{5b}$  are defined as described above, and said intermediates being represented by formula (IV-c), can be prepared by reacting an intermediate of formula (XXXIX) with an appropriate amine, represented by formula (XL), in the presence of an appropriate catalyst, e.g. palladium, and (R)-(+)-2,2'-bis(diphenyl-phosphino)-1,1'-binaphthyl, in a suitable reaction-inert solvent, e.g. tetrahydrofuran.


-26-



- Intermediates of formula (IV) wherein  $\text{R}^{\text{I}}$  is monocyclic heterocycle substituted with  $\text{C}(=\text{O})-\text{NR}^{5a}\text{R}^{5b}$ , wherein  $\text{R}^{5a}$  and  $\text{R}^{5b}$  are defined as described above, said  $\text{R}^{\text{I}}$  being represented by  $\text{R}^{5a}\text{R}^{5b}\text{N}-\text{C}(=\text{O})-\text{R}^{\text{I}'}$ , and said intermediates being represented by
- 5 formula (IV-d), can be prepared by reacting an intermediate of formula (XXXIX) with an appropriate amine, represented by formula (XL), under an atmosphere of carbon monoxide, in the presence of a suitable catalyst, e.g. palladium (II) acetate, and 1,3-bis(diphenylphosphino)propane, in a suitable reaction-inert solvent, e.g. tetrahydrofuran.




- Intermediates of formula (IV) wherein  $\text{P}-\text{Q}_1$  comprises  $\text{C}_{1-10}$ alkyl or  $\text{C}_{3-7}$ cycloalkyl substituted with  $\text{NR}^6-\text{P}$ , said  $\text{C}_{1-10}$ alkyl or  $\text{C}_{3-7}$ cycloalkyl being represented by  $\text{Z}_3$ , said  $\text{P}-\text{Q}_1$  being represented by  $\text{P}-\text{NR}^6-\text{Z}_3-\text{Q}_{1b}$ , and said intermediates being represented by
- 15 formula (IV-e), can be prepared by reacting a compound of formula (I-a-3) with an intermediate of formula (XLI), wherein  $\text{W}_4$  represents a suitable leaving group, such as p-toluenesulphonate. Said reaction can be performed in a reaction-inert solvent, e.g. acetonitrile, in the presence of a suitable base, e.g. dipotassium carbonate.




- Intermediates of formula (IV-e), wherein  $\text{R}^6$  is hydroxy $\text{C}_{1-6}$ alkyl, said intermediates
- 20 being represented by formula (IV-e-1), can be prepared by reacting an intermediate of

-27-

formula (XLII) with an intermediate of formula (XLIII) in the presence of a suitable base, e.g. dipotassium carbonate, and a suitable solvent, e.g. acetonitrile.




Intermediates of formula (XXXIII-a) or (XXXIII-b) can be prepared by protecting an intermediate of formula (XLIV) with a suitable protecting group, such as, for example,  $C_{1-4}alkyloxycarbonyl$ , in a reaction-inert solvent, such as methylene chloride or an alcohol, e.g. methanol, ethanol, 2-propanol and the like, in the presence of a suitable reagent, e.g. di  $C_{1-4}alkyl$  dicarbonate and optionally in the presence of a suitable base, e.g. sodium acetate.




Alternatively, intermediates of formula (XXXIII-a) or (XXXIII-b) can be converted into an intermediate of formula (XLIV) by reaction with a suitable acid, such as hydrochloric acid or hydrobromic acid and the like or mixtures thereof, in the presence of a suitable solvent, e.g. water.

Intermediates of formula (XXXIII-a) or (XXXIII-b), wherein in the definition of  $Q_1$ , the  $X^1$  or  $X^2$  moieties in the radicals of formula (b-1) to (b-8) represent  $NH$ , said  $Q_1$  being represented by  $Q_{1c}-NH$ , and said intermediates by formula (XXXIII-a-1) or (XXXIII-b-1), can be prepared by reacting an intermediate of formula (XLV-a) or (XLV-b), wherein  $W_5$  represents a suitable leaving group, such as for example a halo atom, e.g. chloro, with an intermediate of formula (XLVI).




Intermediates of formula (XLV-a) or (XLV-b) can be prepared by reacting an intermediate of formula (XLVII-a) or (XLVII-b) with  $\text{H}_2\text{P}(=\text{O})(\text{W}_5)_3$  in the presence of a suitable acid, e.g. hydrochloric acid.



5


Intermediates of formula (XLVII-a) or (XLVII-b) can be prepared by reacting an intermediate of formula (XLVIII-a) or (XLVIII-b) with an intermediate of formula (II).




Intermediates of formula (XXXIII-a) can also be prepared by reacting an intermediate of formula (XLVIII-a) with  $\text{P}-\text{Q}_1-\text{C}(=\text{NH})-\text{O}-\text{CH}_2-\text{CH}_3$  in a reaction-inert solvent, such as an alcohol.

10


-29-




Intermediates of formula (XXXV) can be prepared by reacting an intermediate of formula (L) with an intermediate of formula P-Q<sub>1</sub>=C=S, which is synthesized according to the procedures described in EP 0005318, in a reaction-inert solvent, such as an alcohol, e.g. ethanol. To increase the reaction rate, the reaction may be performed at elevated temperatures.




Intermediates of formula (L) can be obtained by reducing an intermediate of formula (LI) in a reaction-inert solvent, e.g. an alcohol, in the presence of a suitable reducing agent, e.g. hydrogen, and an appropriate catalyst, e.g. Raney Nickel.




Intermediates of formula (LI) can be prepared by reacting an intermediate of formula (LII) with an intermediate of formula (LIII), in which W<sub>6</sub> represents a suitable leaving group, such as a halo atom, e.g. chloro. This reaction may be performed in a reaction-inert solvent, e.g. acetonitrile, in the presence of a suitable base, e.g. dipotassium carbonate.




Intermediates of formula (LII) can be prepared by reacting an intermediate of formula (LIV) with a suitable acid, such as hydrochloric acid, in the presence of a suitable solvent, e.g. an alcohol, e.g. ethanol.




- 5 Intermediates of formula (LIV) can be prepared by reacting an intermediate of formula (III) with  $\text{NaN}[\text{C}(=\text{O})\text{H}]_2$ .



- 10 Intermediates of formula (LI) can also be prepared by reacting an intermediate of formula (LV) with an intermediate of formula (LIII) (J. Org. Chem., 25, p 1138, 1960) in a reaction-inert solvent, e.g. *N,N*-dimethylformamide, in the presence of an appropriate base, e.g. sodium hydride.



Intermediates of formula (XXXVI) can be prepared by dehydrating an intermediate of formula (LVI) with a suitable acid, such as sulfuric acid.




15

Intermediates of formula (LVI) wherein, in the definition of  $\text{Q}_{1a}$ , the  $\text{X}^1$  or  $\text{X}^2$  moieties are  $\text{CH}_2$ , said  $\text{Q}_{1a}$  being represented by  $\text{Q}_{1a'}$ , and said intermediates being represented by formula (LVI-a), can be prepared by reacting a carbonyl moiety of formula (LVII) with an intermediate of formula (LVIII) in the presence of *N,N*-diisopropylamine and butyl lithium, in a suitable reaction-inert solvent, e.g. tetrahydrofuran.


20




-31-




Intermediates of formula (IV), wherein G is C<sub>1-10</sub>alkanediyl substituted with C<sub>1-6</sub>alkyloxy, arylC<sub>1-6</sub>alkyloxy, HO(-CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-, C<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-, or arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>CH<sub>2</sub>O)<sub>n</sub>-, said group of substituents being represented by O-Z<sub>4</sub>, said G being represented by Z<sub>4</sub>-O-G<sub>1</sub>, and said intermediates being represented by formula (IV-f), can be prepared by reacting an intermediate of formula (XXXIII-a), with an intermediate of formula (LIX), optionally in the presence of a suitable acid, such as p-toluenesulfonic acid and the like, and optionally in the presence of a suitable solvent, such as *N,N*-dimethylacetamide. To increase the reaction rate, the reaction may be carried out at elevated temperatures.



Intermediates of formula (LIX) can be prepared by reacting an intermediate of formula (LX) with a reagent of formula (LXI) or (LXII) in a reaction-inert solvent, such as an alcohol, or toluene, in the presence of an acid, e.g. 4-methylbenzenesulphonic acid.




Intermediates of formula (LX) can be prepared by oxidizing an intermediate of formula (LXIII) with a suitable oxidizing agent, e.g. MnO<sub>2</sub>, in a reaction-inert solvent, such as methylene chloride.




Intermediates of formula (IV-f) can also be prepared by reacting an intermediate of formula (IV) wherein G is C<sub>1-10</sub>alkanediyl substituted with hydroxy, said G being represented by G<sub>1</sub>-OH, and said intermediates being represented by formula (IV-g),

with an intermediate of formula (LXIV), wherein  $W_7$  is a suitable leaving group, such as a halo atom, e.g. iodo, in the presence of a suitable base, e.g. sodium hydride, in a reaction-inert solvent, e.g. tetrahydrofuran.




- 5 Intermediates of formula (IV-g), wherein the carbon atom of  $G_1$  carrying the hydroxy, also carries a hydrogen atom, said  $G_1\text{-OH}$  being represented by  $H\text{-}G_2\text{-OH}$ , and said intermediates being represented by formula (IV-g-1), can be prepared by reducing an intermediate of formula (LXV) in the presence of a suitable reducing agent, e.g. sodium borohydride, in a reaction-inert solvent, such as an alcohol, tetrahydrofuran or a mixture thereof. Intermediates of formula (LXV) can also first be deprotected, e.g. in the presence of a suitable acid, such as hydrochloric acid and the like, resulting in intermediates of formula (LXVI), followed by a reduction, resulting in a compound of formula (I-q-1) wherein  $Q$  represents  $H\text{-}Q_1$ , said compounds being represented by formula (I-q-1-1).
- 10




15

Intermediates of formula (IV), wherein  $G$  is ethyl substituted with hydroxy, said intermediates being represented by formula (IV-g-2) can also be prepared by reacting an


intermediate of formula (XXXIII-a) with an intermediate of formula (LXVII) in the presence of a suitable base, such as sodium hydride, in a reaction-inert solvent, such as *N,N*-dimethylformamide.




- 5 A subgroup of intermediates of formula (IV-g-2), represented by formula (IV-g-2-1), can also be prepared by reacting an intermediate of formula (LXVIII) with an intermediate of formula (LXIX) in the presence of 1,3-dicyclohexylcarbodiimide, in a reaction-inert solvent, e.g. toluene.




- 10 Intermediates of formula (LXV) can be prepared by reacting an intermediate of formula (XXXIII-a) with an intermediate of formula (LXX), wherein W<sub>8</sub> is a suitable leaving group, such as a halo atom, e.g. bromo, in the presence of a suitable base, e.g. sodium hydride, in a reaction-inert solvent, e.g. *N,N*-dimethylformamide.




Intermediates of formula (V) can be prepared by reacting an intermediate of formula (LXXI) with 1*H*-isoindole-1,3 (2*H*)-dione in the presence of triphenylphosphine and diethyl azodicarboxylate.



- 5 Intermediates of formula (V) may also be prepared by reacting an intermediate of formula (LXXII) with 1*H*-isoindole-1,3 (2*H*)-dione in the presence of a suitable base, such as sodium hydride, and a suitable solvent, such as *N,N*-dimethylformamide.




- 10 Intermediates of formula (LXXII) can be prepared by reacting an intermediate of formula (LXXI) with an intermediate of formula (LXXIII), wherein  $W_9$  represents a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of a suitable base, such as *N,N*-diethyl-ethanamine, and a suitable solvent, such as methylene chloride.




- 15 Intermediates of formula (V), wherein in the definition of  $Q_2$ ,  $R^2$  is  $C_{1-10}$ alkyl, said  $Q_2$  being represented by  $C_{1-10}$ alkyl- $Q_{1b}$ , and said intermediates by formula (V-a), can be prepared by reacting a compound of formula (I-a-3) with an intermediate of formula (LXXIV), wherein  $W_{10}$  is a suitable leaving group, such as a halo atom, e.g. chloro, in

the presence of a suitable base, such as dipotassium carbonate, and a suitable solvent, such as acetonitrile.




Intermediates of formula (LXXI) wherein, in the definition of Q<sub>2</sub>, the carbon atom  
 5 carrying the hydroxy, also carries two hydrogen atoms, said HO-Q<sub>2</sub> being represented  
 by HO-CH<sub>2</sub>-Q<sub>2</sub>, and said intermediates being represented by formula (LXXI-a), can be  
 prepared by reducing an intermediate of formula (LXXV) in the presence of a suitable  
 reducing agent, such as lithium aluminium hydride, in a suitable reaction-inert solvent,  
 e.g. tetrahydrofuran.




10


Intermediates of formula (LXXI), wherein, in the definition of Q<sub>2</sub>, the carbon atom  
 carrying the hydroxy, carries also at least one hydrogen, said HO-Q<sub>2</sub> being represented  
 by HO-Q<sub>3</sub>H, and said intermediates being represented by formula (LXXI-b), can be  
 prepared by reducing an intermediate of formula (IX) with a suitable reducing agent,  
 15 e.g. sodium borohydride, in a reaction-inert solvent, e.g. an alcohol.




Intermediates of formula (VI) wherein, in the definition of  $Q_2$ ,  $R^2$  is  $C_{1-10}$ alkyl substituted with  $N(P)_2$  and the carbon atom adjacent to the nitrogen atom carrying the  $R^2$  substituent carries also at least one hydrogen atom, said  $Q_2$  being represented by  $(P)_2N-C_{1-10}alkyl-NH-Q_{2a}H$ , and said intermediates being represented by formula (VI-a),  
 5 can be prepared by reductive amination of an intermediate of formula (LXXVI) with an intermediate of formula (LXXVII) in the presence of a suitable reductive agent, such as hydrogen, and a suitable catalyst, such as palladium-on-charcoal, platinum-on-charcoal, and the like, and optionally in the presence of a suitable catalyst poison, such as a thiophene solution. A suitable solvent in this reaction is a reaction-inert solvent, such  
 10 as an alcohol.




Intermediates of formula (LXXVI) can be prepared by deprotecting an intermediate of formula (LXXVIII) in the presence of a suitable acid, such as hydrochloric acid and the like, in a suitable solvent, e.g. water.




15 Intermediates of formula (IX) may be prepared by deprotecting an intermediate of formula (LXXIX) in the presence of a suitable acid, e.g. hydrochloric acid and the like.




20 Intermediates of formula (LXXIX) can be prepared by reacting an intermediate of formula (LXXX) with an intermediate of formula (III) in the presence of a suitable base, e.g. dipotassium carbonate, in a suitable reaction-inert solvent, e.g. acetonitrile.




- Intermediates of formula (LXXX) wherein, in the definition of  $Q_3$ , the  $X^1$  or  $X^2$  moiety of the radicals of formula (b-1) to (b-8) represent NH, said  $Q_3$  being represented by  $Q_3-NH$ , and said intermediates being represented by formula (LXXX-a), may be prepared by cyclizing an intermediate of formula (LXXXI) in the presence of mercury oxide and sulphur, in a suitable reaction-inert solvent, e.g. an alcohol.



- Intermediates of formula (LXXXI) can be prepared by reducing an intermediate of formula (LXXXII) in the presence of a suitable reducing agent, such as hydrogen, in the presence of a suitable catalyst, such as palladium-on-charcoal, platinum-on-charcoal and the like, in a suitable solvent, e.g. a mixture of ammonia in alcohol. Suitable alcohols are methanol, ethanol, 2-propanol and the like.




- Intermediates of formula (LXXXII) can be prepared by reacting an intermediate of formula (LXXXIII) with an intermediate of formula (LXXXIV) in a suitable reaction-inert solvent, e.g. ethanol.




Intermediates of formula (IX), wherein, in the definition of  $Q_3$ ,  $R^2$  comprises  $C_{1-10}$ alkyl, said  $Q_3$  being represented by  $C_{1-10}$ alkyl- $Q_{1b}$ , and said intermediates being represented


- by formula (IX-a), can be prepared by reacting a compound of formula (I-a-3) with a reagent of formula (LXXXV), wherein  $(O=)C_{1-10}alkyl$  represents a carbonyl derivative of  $C_{1-10}alkyl$  and wherein  $W_{11}$  is a suitable leaving group, such as a halo atom, e.g. bromo, in a reaction-inert solvent, e.g. acetonitrile, in the presence of a suitable base,
- 5 e.g. dipotassium carbonate.



- Intermediates of formula (X) wherein  $Q_4$  comprises  $C_{1-9}alkyl$ , said  $Q_4$  being represented by  $C_{1-9}alkyl-Q_{1b}$ , and said intermediates being represented by formula (X-a), can be prepared by reacting a compound of formula (I-a-3) with a reagent of formula
- 10 (LXXXVI) wherein  $W_{12}$  represents a suitable leaving group, such as a halo atom, e.g. chloro, in a reaction-inert solvent, e.g. 3-methyl-2-butanone, in the presence of a suitable base, e.g. dipotassium carbonate, sodium bicarbonate and the like.




- Intermediates of formula (X), wherein  $NC-Q_4$  represents  $NC-(C_{1-9}alkyl)(R^4)N-C(=O)-$
- 15  $Alk-X^1$ , said intermediates being represented by formula (X-b), can be prepared by reacting an intermediate of formula (LXXXVII) with an intermediate of formula (LXXXVIII) in the presence of di-1H-imidazol-2-yl-methanone, a suitable base, such as *N,N*-diethyl-ethanamine, and a suitable solvent, such as methylene chloride.






Intermediates of formula (XI), wherein  $Q_4$  represents  $Q_{1b}$ , said intermediates being represented by formula (XI-a), can be prepared by reacting a compound of formula (I-a-3) with an intermediate of formula (LXXXIX), wherein  $W_{13}$  represents a suitable leaving group, such as a halo atom, e.g. chloro, in the presence of a suitable base, such as disodium carbonate, and in the presence of a suitable solvent, such as 3-methyl-2-butanone.



Intermediates of formula (XIX) can be prepared by reacting an intermediate of formula (XC) with a suitable acid, such as hydrochloric acid.



10

Pure stereochemically isomeric forms of the compounds of formula (I) may be obtained by the application of art-known procedures. Diastereomers may be separated by physical methods such as selective crystallization and chromatographic techniques, e.g., counter-current distribution, liquid chromatography and the like.

The compounds of formula (I) as prepared in the hereinabove described processes are generally racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) which are sufficiently basic or acidic may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid, respectively chiral base. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali or acid. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase. Said pure stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the

20

25

-40-

reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods will advantageously employ enantiomerically pure starting materials.

- 5 The compounds of formula (I) show antiviral properties. Viral infections treatable using the compounds and methods of the present invention include those infections brought on by ortho- and paramyxoviruses and in particular by human and bovine respiratory syncytial virus (RSV).
- 10 The *in vitro* antiviral activity against RSV of the present compounds was tested in a test as described in the experimental part of the description, and may also be demonstrated in a virus yield reduction assay. The *in vivo* antiviral activity against RSV of the present compounds may be demonstrated in a test model using cotton rats as described in Wyde et al. (Antiviral Research (1998), 38, 31-42).

15

- Due to their antiviral properties, particularly their anti-RSV properties, the compounds of formula (I) or any subgroup thereof, their prodrugs, *N*-oxides, addition salts, quaternary amines, metal complexes and stereochemically isomeric forms, are useful in the treatment of individuals experiencing a viral infection, particularly a RSV infection,
- 20 and for the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses, in particular the respiratory syncytial virus.

- The compounds of the present invention or any subgroup thereof may therefore be used
- 25 as medicines. Said use as a medicine or method of treatment comprises the systemic administration to viral infected subjects or to subjects susceptible to viral infections of an amount effective to combat the conditions associated with the viral infection, in particular the RSV infection.

- 30 The present invention also relates to the use of the present compounds or any subgroup thereof in the manufacture of a medicament for the treatment or the prevention of viral infections, particularly RSV infection.

- The compounds of the present invention or any subgroup thereof may be formulated
- 35 into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form or metal

complex, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for  
5 administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, lubricants, binders,  
10 disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid  
15 solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before  
20 use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin.

25 The compounds of the present invention may also be administered via oral inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder, a solution being preferred. Any system developed for the delivery of solutions,  
30 suspensions or dry powders via oral inhalation or insufflation are suitable for the administration of the present compounds.

Thus, the present invention also provides a pharmaceutical composition adapted for administration by inhalation or insufflation through the mouth comprising a compound of formula (I) and a pharmaceutically acceptable carrier. Preferably, the compounds of  
35 the present invention are administered via inhalation of a solution in nebulized or aerosolized doses.

It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, suppositories, powder packets, wafers, injectable solutions or suspensions and the like, and segregated multiples thereof.

In general it is contemplated that an antivirally effective daily amount would be from 0.01 mg/kg to 500 mg/kg body weight, more preferably from 0.1 mg/kg to 50 mg/kg body weight. It may be appropriate to administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight, sex, extent of disorder and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines.

Also, the combination of another antiviral agent and a compound of formula (I) can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of formula (I), and (b) another antiviral compound, as a combined preparation for simultaneous, separate or sequential use in antiviral treatment. The different drugs may be combined in a single preparation together with pharmaceutically acceptable carriers. For instance, the compounds of the present invention may be combined with interferon-beta or tumor necrosis factor-alpha in order to treat or prevent RSV infections.

The following examples are intended to illustrate the present invention.

### A. Preparation of the intermediate compounds

#### Example A1

- 5 a)  $K_2CO_3$  (0.129 mol) was suspended in a solution of ethyl 4-(1*H*-benzimidazol-2-yl-amino)-1-piperidinecarboxylate (0.0347 mol) and 2-bromo-1-(4-chlorophenyl)ethanone (0.0647 mol) in acetonitrile (150ml). The mixture was stirred and refluxed for 8 hours, then cooled, poured out into  $H_2O$  and extracted with  $CH_2Cl_2$ . The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $CH_2Cl_2/CH_3OH/NH_4OH$  97.5/2.5/0.1). The pure fractions were collected and the solvent was evaporated. Part of this fraction
- 10 (3g) was taken up in 2-propanone and diethyl ether. The precipitate was filtered off and dried, yielding 2g of ethyl 4-[[1-[2-(4-chlorophenyl)-2-oxoethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (interm. 1).
- b) A mixture of intermediate (1) (0.015 mol) in  $HCl$  12N (100ml) was stirred and refluxed for 12 hours, then the solvent was evaporated. Ethylacetate was added. The mixture was basified with a saturated  $NaHCO_3$  solution. The precipitate was filtered
- 15 off, washed with  $H_2O$  and with ethylacetate and dried. The residue (5.5g) was crystallized from ethylacetate. The precipitate was filtered off and dried, yielding 4.8g of 1-(4-chlorophenyl)-2-[2-(4-piperidinylamino)-1*H*-benzimidazol-1-yl]ethanone dihydrate (80%) (interm. 2).

#### 20 Example A2

- a)  $NaBH_4$  (0.034 mol) was added portionwise at 5°C to a mixture of (±)-1,1-dimethylethyl 4-[[1-(2-oxo-2-phenylethyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (0.034 mol) in tetrahydrofuran (250ml) and methanol (250ml). The mixture was stirred at 5°C and then hydrolyzed cold with  $H_2O$ . The solvent was evaporated and the
- 25 residue was taken up in  $H_2O$ . The precipitate was filtered off, washed with diisopropyl ether and dried, yielding 11.3g of (±)-1,1-dimethylethyl 4-[[1-(2-hydroxy-2-phenylethyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (76%) (interm. 3).
- b) A mixture of intermediate (3) (0.0183 mol) in tetrahydrofuran (50ml) was cooled to 0°C under  $N_2$  flow.  $NaH$  80% (0.0366 mol) was added portionwise. The mixture was
- 30 brought to room temperature, then stirred at room temperature for 30 minutes and cooled again to 0°C. A solution of  $CH_3I$  (0.0183 mol) in tetrahydrofuran (50ml) was added dropwise. The mixture was stirred at room temperature for 2 hours, then cooled, hydrolyzed and extracted with ethylacetate. The organic layer was separated, washed with  $H_2O$ , dried, filtered and the solvent was evaporated. The residue was purified by
- 35 column chromatography over silica gel (eluent:  $CH_2Cl_2/CH_3OH/NH_4OH$  98.5/1.5/0.1). The desired fractions were collected and the solvent was evaporated, yielding 5g of

(±)-1,1-dimethylethyl 4-[[1-(2-methoxy-2-phenylethyl)-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (interm. 4).

### Example A3

a) NaOCH<sub>3</sub> (0.2 mol) was added to a mixture of *N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine dihydrobromide (0.1 mol) in methanol (389ml), the mixture was cooled on an ice bath and stirred for 2 hours. Bis(1,1-dimethylethyl) dicarbonoate (0.1mol) was added to a cooled mixture on an ice bath and then stirred for 18 hours at room temperature. The solvent was evaporated and the residue suspended in water/diisopropyl ether. The residue was filtered off, washed with water/diisopropyl ether and dried. The residue was boiled up in CH<sub>3</sub>OH, yielding 17.46g of 1,1-dimethylethyl 4-(1*H*-benzimidazol-2-ylamino)-1-piperidinecarboxylate (55.2%) (interm. 5).

b) A mixture of 3-(benzyloxy)-6-methyl-2-pyridinemethanol (0.0314 mol) and MnO<sub>2</sub> (29.52g) in CH<sub>2</sub>Cl<sub>2</sub> (100ml) was stirred at room temperature overnight and then purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub> 100%). The pure fractions were collected and the solvent was evaporated, yielding 6.71g of 6-methyl-3-(phenylmethoxy)-2-pyridinecarboxaldehyde (94%) (interm. 6).

c) A mixture of intermediate (6) (0.0385 mol) and triethylorthoformate in the presence of 4-methylbenzenesulfonic acid (0.5g) in toluene (200ml) was stirred and refluxed for 6 hours. The solvent was evaporated. The residue was taken up in H<sub>2</sub>O, Na<sub>2</sub>CO<sub>3</sub> and CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 9.6g of 2-(diethoxymethyl)-6-methyl-3-(phenylmethoxy)pyridine (83%) (interm. 7).

d) Intermediate (7) (0.03185 mol) and intermediate (5) (0.03185 mol) were heated to 150°C and purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 98/2). The pure fractions were collected and the solvent was evaporated, yielding 10.25g of (±)-1,1-dimethylethyl 4-[[1-[ethoxy[6-methyl-3-(phenylmethoxy)-2-pyridinyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (56%) (interm. 8).

e) A mixture of 2-(diethoxymethyl)-6-bromo-pyridine (0.03 mol), intermediate 5 (0.03 mol) and 4-methylbenzenesulfonic acid (2g) in toluene (700ml) was stirred and refluxed for 20 hours using a water separator. 4-Methylbenzenesulfonic acid was added and the mixture was stirred and refluxed for 48 hours. 4-Methylbenzenesulfonic acid was added again and the mixture was stirred and refluxed for another 48 hours. 4-Methylbenzenesulfonic acid was added again. The mixture was stirred and refluxed for 24 hours, then cooled and washed with a diluted NaOH solution. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by

column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/C<sub>2</sub>H<sub>5</sub>OH 95/5). The pure fractions were collected and the solvent was evaporated. The residue was suspended in petroleum ether. The precipitate was filtered off and dried, yielding 1.4g of (±)-1,1-dimethylethyl 4-[[1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (9%) (interm. 33).

#### Example A4

- a) A mixture of 2,3-pyridinediamine (0.05 mol) and ethyl 4-(2-ethoxy-2-iminoethyl)-1-piperidinecarboxylate monohydrochloride (0.05 mol) in methanol (150ml) was stirred and refluxed for 3 days. The solvent was evaporated and the residue was taken up in CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with K<sub>2</sub>CO<sub>3</sub> 10%, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 94/6/0.1). The pure fractions were collected and the solvent was evaporated, yielding 7.6g of ethyl 4-[(1*H*-imidazo[4,5-*b*]pyridin-2-yl)methyl]-1-piperidinecarboxylate (52%) (interm. 9).
- b) NaH (0.028 mol) was added portionwise at 0°C to a mixture of intermediate (9) (0.023 mol) in *N,N*-dimethylformamide (75ml). 2-Bromo-1-phenylethanone (0.028 mol) was added. The mixture was stirred at room temperature for 1 hour. H<sub>2</sub>O was added and the mixture was extracted with ethylacetate. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 97.5/2.5/0.1). The pure fractions were collected and the solvent was evaporated, yielding 4.7g of ethyl 4-[[1-(2-oxo-2-phenylethyl)-1*H*-imidazo[4,5-*b*]pyridin-2-yl)methyl]-1-piperidinecarboxylate (50.5%) (interm. 10).
- c) NaBH<sub>4</sub> (0.0137 mol) was added portionwise at 5°C under N<sub>2</sub> flow to a mixture of intermediate (10) (0.0137 mol) in methanol (100ml). The mixture was hydrolyzed with H<sub>2</sub>O and extracted with ethylacetate. The organic layer was separated, dried, filtered and the solvent was evaporated, yielding 5.6g of (±)-ethyl 4-[[1-(2-hydroxy-2-phenylethyl)-1*H*-imidazo[4,5-*b*]pyridin-2-yl)methyl]-1-piperidinecarboxylate (interm. 11).

#### Example A5

- A mixture of (±)-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine (0.00205 mol), 1-chloro-2-propanone (0.00308 mol) and K<sub>2</sub>CO<sub>3</sub> (0.0041 mol) in acetonitrile (8ml) was stirred and refluxed for 8 hours. H<sub>2</sub>O was added and the mixture was extracted with ethylacetate. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/ NH<sub>4</sub>OH 97/3/0.1). The pure fractions were collected and the solvent was evaporated, yielding: 0.67g of (±)-1-[4-[[1-

[ethoxy(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]-2-propanone (77%) (interm. 12).

#### Example A6

4-Methylbenzenesulfonyl chloride (0.2222 mol) was added portionwise at 10°C to a mixture of 1,1-dimethylethyl [1-(hydroxymethyl)-2-methylpropyl]carbamate (0.202 mol) in pyridine (65ml). The mixture was stirred at 10°C for 2 hours. H<sub>2</sub>O (75ml) was added at 10°C. The precipitate was filtered off, washed with H<sub>2</sub>O and taken up in CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with H<sub>2</sub>O, dried, filtered and the solvent was evaporated, yielding 49g of (±)-1,1-dimethylethyl [1-[[[(4-methylphenyl)sulfonyl]oxy]methyl]-2-methylpropyl]carbamate (68%) (interm. 13).

#### Example A7

a) A mixture of (±)-1,1-dimethylethyl 4-[[1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (0.00189 mol) (interm. 33), Pd (0.026g), (R)-(+)-2,2'-bis(diphenyl-phosphino)-1,1'-binaphthyl (0.046g) and NH(CH<sub>3</sub>)<sub>2</sub> gas (10g) in tetrahydrofuran (200ml) was stirred in an autoclave at 100°C for 16 hours under pressure of CO (30 atm). The mixture was filtered and the filtrate was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/ NH<sub>3</sub>) 99/1). The pure fractions were collected and the solvents was evaporated, yielding 0.8g of (±)-1,1-dimethylethyl 4-[[1-[[6-(dimethylamino)-2-pyridinyl]ethoxymethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (86%) (interm. 14).

b) A mixture of intermediate 33 (0.0032 mol), Pd(OAc)<sub>2</sub> (0.030 g) and 1,3-propanediylbis[diphenylphosphine] (0.110 g) in tetrahydrofuran (100 ml) under ammonia (liq., 10 atm) and CO (gas, 30 atm) was stirred for 16 hours at 100 °C. The solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 98/2). The pure fractions were collected and the solvent was evaporated, yielding 0.15 g of interm. 41.

#### Example A8

A mixture of α-[[[(3-amino-2-pyridinyl)amino]methyl]benzenemethanol (0.043 mol) and ethyl 4-isothiocyanato-1-piperidinecarboxylate (0.047 mol) in toluene (200ml) was stirred and refluxed for 30 minutes. *N,N*-methanetetrayl-biscyclohexanamine (0.065 mol) was added and the mixture was stirred and refluxed overnight. The solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 96/4/0.2). The pure fractions were collected and the solvent was evaporated. Part of the residue (1.5g) was crystallized from diisopropyl ether. The precipitate was filtered off and dried, yielding 1.35g of (±)-ethyl 4-[[1-(2-



hydroxy-2-phenylethyl)-1*H*-imidazo[4,5-*b*]pyridin-2-yl]amino]-1-piperidinecarboxylate (interm. 15).


#### Example A9

Reaction under N<sub>2</sub> flow. NaH 60% (0.02 mol) was added to a mixture of 1,1-dimethylethyl 4-(1*H*-benzimidazol-2-ylamino)-1-piperidinecarboxylate (0.02 mol) in *N,N*-dimethylformamide (100ml). The mixture was stirred at 40°C for 1 hour.

6-(Epoxyethyl)-2-picoline (0.02 mol) in a small amount of *N,N*-dimethylformamide was added. The mixture was stirred at 100°C overnight. The solvent was evaporated. The residue was taken up in H<sub>2</sub>O and CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5 and 90/10). The pure fractions were collected and the solvent was evaporated, yielding 3.5g of (±)-1,1-dimethylethyl 4-[[1-[2-hydroxy-2-(6-methyl-2-pyridinyl)ethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (interm. 16).


Tables 1,2 and 3 list intermediates which were prepared analogous to one of the above examples.

Table 1




Int. No.	Ex. No.	L	R <sup>a</sup>	Physical data
17	A1b	H	2-Cl	H <sub>2</sub> O (1:1)
18	A1b	H	4-OCH <sub>3</sub>	
19	A1b	H	3-Cl	
20	A1b	H	3-F	
2	A1b	H	4-Cl	
21	A1b	H	3-CH <sub>3</sub>	H <sub>2</sub> O (1:2)
22	A1b	H	2-CH <sub>3</sub>	
23	A1a	-C(=O)-O-C <sub>2</sub> H <sub>5</sub>	4-CH <sub>3</sub>	H <sub>2</sub> O (1:1)
24	A1b	H	4-CH <sub>3</sub>	H <sub>2</sub> O (1:1); mp. 180°C
25	A1b	H	3-OCH <sub>3</sub>	H <sub>2</sub> O (1:1), HCl (1:1); mp. 220°C
26	A1b	H	2-F	

Table 2



Int. No.	Ex. No.	L	n	a	b	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	Physical data; mp.
27	A3e	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	85°C
28	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	CH <sub>3</sub>		H	
29	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>2</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
30	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	CH <sub>3</sub>	-[(CH <sub>2</sub> ) <sub>2</sub> -O] <sub>2</sub> -CH <sub>3</sub>	H	
31	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	Phenyl	-(CH <sub>2</sub> ) <sub>2</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
32	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	-CH <sub>2</sub> -O-CH <sub>3</sub>	-CH <sub>3</sub>	H	
33	A3e	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	Br	-C <sub>2</sub> H <sub>5</sub>	H	
34	A3d	-CH <sub>2</sub> -phenyl	0	N	CH	H	-C <sub>2</sub> H <sub>5</sub>	H	
35	A5	-CH <sub>2</sub> -C(=O)-CH(CH <sub>3</sub> ) <sub>2</sub>	0	N	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
36	A5	-CH <sub>2</sub> -C(=O)-CH(CH <sub>3</sub> ) <sub>2</sub>	0	N	CH	CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>2</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
37	A5	-CH <sub>2</sub> -C(=O)-CH(CH <sub>3</sub> ) <sub>2</sub>	0	N	CH	CH <sub>3</sub>	-[(CH <sub>2</sub> ) <sub>2</sub> -O] <sub>2</sub> -CH <sub>3</sub>	H	
38	A5	-CH <sub>2</sub> -C(=O)-CH(CH <sub>3</sub> ) <sub>2</sub>	0	N	CH	Phenyl	-(CH <sub>2</sub> ) <sub>2</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
40	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	3-O-benzyl	
41	A7b	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	-CO-NH <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
42	A7b	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	-CO-N(CH <sub>3</sub> ) <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
16	A9	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	1	N	CH	CH <sub>3</sub>	H	H	
44	A3d	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>2</sub> -OCH <sub>3</sub>	H	
4	A2b	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	1	CH	CH	H	CH <sub>3</sub>	H	
15	A8	-C(=O)-O-C <sub>2</sub> H <sub>5</sub>	1	CH	N	H	H	H	
47	A7a	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	0	N	CH	-N(CH <sub>3</sub> ) <sub>2</sub>	C <sub>2</sub> H <sub>5</sub>	H	

Table 3



Int. No.	Ex. No.	b	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	L	Physical data; mp.
48	A3d	CH	Br	-CH <sub>2</sub> -CH <sub>3</sub>	H	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	
49	A5	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	Cl	-CH <sub>2</sub> -C(=O)-CH(CH <sub>3</sub> ) <sub>2</sub>	
50	A3d	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	Cl	-C(=O)-O-C(CH <sub>3</sub> ) <sub>3</sub>	
51	A5	N	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	H	-CH <sub>2</sub> -C(=O)-CH(CH <sub>3</sub> ) <sub>2</sub>	
52	A3d	N	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	H	-CH <sub>2</sub> -C <sub>6</sub> H <sub>5</sub>	

### B. Preparation of the final compounds

#### 5 Example B1

A mixture of intermediate (4) (0.0102 mol) in HCl 3N (80ml) and 2-propanol (10ml) was stirred at 40°C for 2 hours. The mixture was brought to room temperature and poured out on ice. CH<sub>2</sub>Cl<sub>2</sub> was added. The mixture was basified with K<sub>2</sub>CO<sub>3</sub> solid, stirred at room temperature for 1 hour and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was separated, washed with H<sub>2</sub>O, dried, filtered and the solvent was evaporated. The residue was crystallized from diethyl ether and CH<sub>3</sub>OH. The precipitate was filtered off and dried, yielding 2.9g of (±)-1-(2-methoxy-2-phenylethyl)-N-(4-piperidiny)-1H-benzimidazol-2-amine (81%) (compound 1).

#### Example B2

A mixture of intermediate (11) (0.0139 mol) and KOH (0.1 mol) in 2-propanol (200ml) was stirred and refluxed overnight. The solvent was evaporated and the residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 80/20/3). The pure fractions were collected and the solvent was evaporated. The residue was converted into the ethanedioic acid salt (1:2) with ethanedioic acid. The mixture was crystallized from 2-propanone. The precipitate was filtered off and dried, yielding 3.9g of (±)-α-phenyl-2-(4-piperidinylmethyl)-1H-imidazo[4,5-b]pyridine-1-ethanol ethanedioate (1:2) (compound 2).

Example B3

- a) A mixture of intermediate (8) (0.00175 mol) in trifluoroacetic acid (20ml) and  $\text{CH}_2\text{Cl}_2$  (50ml) was stirred at room temperature for 2 hours, poured out into ice water and alkalized with a NaOH solution.  $\text{CH}_2\text{Cl}_2$  was added. The organic layer was  
5 separated, dried, filtered and the solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  90/10). The pure fractions were collected and the solvent was evaporated. The residue was converted into the hydrochloric acid salt (1:3). The precipitate was filtered off and dried, yielding 0.48g of ( $\pm$ )-1-[ethoxy[6-methyl-(3-phenylmethoxy)-2-pyridinyl]methyl]-*N*-(4-piperidinyl)-  
10 1*H*-benzimidazol-2-amine trihydrochloride dihydrate 2-propanolate (1:1) (compound 3).
- b) A mixture of ( $\pm$ )-1,1-dimethylethyl 4-[[1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinecarboxylate (0.0026 mol) in 2-propanol (30ml) and  $\text{HBr}/\text{CH}_3\text{COOH}$  (2ml) was stirred and refluxed for 2 hours and then cooled. The  
15 solvent was evaporated. The residue was taken up in  $\text{H}_2\text{O}$  and  $\text{CH}_2\text{Cl}_2$ . The mixture was alkalized with a NaOH solution. The organic layer was separated, washed with  $\text{H}_2\text{O}$ , dried, filtered and the solvent evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/\text{NH}_3$  90/10). The pure fractions were collected and the solvent was evaporated. The residue was suspended in petroleum  
20 ether. The precipitate was filtered off and dried. This fraction was recrystallized from a small amount of  $\text{CH}_3\text{CN}$ . The precipitate was filtered off and dried, yielding 0.22g of ( $\pm$ )-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-*N*-(4-piperidinyl)-1*H*-benzimidazol-2-amine (19.6%) (compound 4).

Example B4

- 25 A mixture of ( $\pm$ )-1-[ethoxy(2-pyridinyl)methyl]-*N*-[1-(phenylmethyl)-4-piperidinyl]-1*H*-benzimidazol-2-amine (0.011 mol) in methanol (150ml) was hydrogenated for 4 days with Pd/C 10% (2g) as a catalyst. After uptake of  $\text{H}_2$  (1 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  90/10). The pure  
30 fractions were collected and the solvent was evaporated, yielding 1.5g of ( $\pm$ )-1-[ethoxy-(2-pyridinyl)methyl]-*N*-(4-piperidinyl)-2-benzimidazol-2-amine (39%) (compound 5).

Example B5

- $\text{NaBH}_4$  (0.0078 mol) was added portionwise to a mixture of intermediate (2) (0.0078 mol) in tetrahydrofuran (50ml) and methanol (50ml), and the mixture was stirred at 5°C  
35 under  $\text{N}_2$  flow for 2 hours. The mixture was hydrolyzed cold with  $\text{H}_2\text{O}$  (3ml) and the solvent was evaporated. The precipitate was filtered off, washed with  $\text{H}_2\text{O}$  and dried.

The residue (3g) was crystallized from diisopropyl ether. The precipitate was filtered off and dried, yielding 2.9g of ( $\pm$ )- $\alpha$ -(4-chlorophenyl)-2-(4-piperidinylamino)-1*H*-benzimidazole-1-ethanol (100%) (compound 6).

#### Example B6

- 5 A mixture of compound (4) (0.0035 mol), 1,1-dimethylethyl (2-bromomethyl)-carbamate (0.005 mol) and Na<sub>2</sub>CO<sub>3</sub> (0.01 mol) in 2-butanone (100ml) was stirred and refluxed for 20 hours. H<sub>2</sub>O was added. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH 95/5 to 90/10). The pure fractions were  
10 collected and the solvent was evaporated, yielding 1.3g of ( $\pm$ )-1,1-dimethylethyl [2-[4-[[1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]-ethyl]carbamate (compound 7).

#### Example B7

- A mixture of compound (4) (0.00348 mol), intermediate (13) (0.00348 mol) and K<sub>2</sub>CO<sub>3</sub>  
15 (0.01392 mol) in acetonitrile (20ml) and *N,N*-dimethylformamide (4ml) was stirred at 60°C for 4 hours (1 equivalent of intermediate (13) was added every hour) and then cooled. The solvent was evaporated. The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with H<sub>2</sub>O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent:  
20 CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 96.5/3.5/0.1). Two pure fractions were collected and their solvents were evaporated, yielding 1g of ( $\pm$ )-1,1-dimethylethyl [1-[4-[[1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]methyl]-2-methylpropyl]carbamate (47%) (compound 8).

#### Example B8

- 25 A mixture of compound (7) (0.0026 mol) in 2-propanol (30ml) and HBr/acetic acid (2ml) was stirred and refluxed for 90 minutes and then cooled. The solvent was evaporated. The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub> and H<sub>2</sub>O. The organic layer was separated, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 90/10). The  
30 pure fractions were collected and the solvent was evaporated. The residue was suspended in diisopropyl ether. The precipitate was filtered off and dried. This fraction was purified again by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 90/10). The pure fractions were collected and the solvent was evaporated. The residue was suspended in diisopropyl ether. The precipitate was  
35 filtered off and dried, yielding 0.23g of ( $\pm$ )-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-amine (compound 9).

Example B9

A mixture of compound (8) (0.00162 mol) in 2-propanol/HCl (1ml) and 2-propanol (10ml) was stirred and refluxed for 1 hour and then cooled. The solvent was evaporated. The residue was taken up in CH<sub>2</sub>Cl<sub>2</sub>. The organic solution was washed with K<sub>2</sub>CO<sub>3</sub> 10% and with H<sub>2</sub>O, dried, filtered and the solvent was evaporated. The residue was purified by column chromatography (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 94/6/1). The pure fractions were collected and the solvent was evaporated, yielding 0.23g of (±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidiny]-1-[(6-bromo-2-pyridinyl)-ethoxymethyl]-1*H*-benzimidazol-2-amine (27%) (compound 10).

10 Example B10

A mixture of (±)-1,1-dimethylethyl [2-[4-[[1-[ethoxy[6-methyl-3-(phenylmethoxy)-2-pyridinyl]methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidinyl]ethyl]carbamate (0.0016 mol) and KOH (1g) in sec-butanol (25ml) was stirred and refluxed for 6 hours. The solvent was evaporated. The residue was purified over silica gel on a glass filter (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 95/5, 93/7 to 90/10). The pure fractions were collected and the solvent was evaporated. The residue was converted into the hydrochloric acid salt (1:3). The precipitate was filtered off and dried, yielding 0.5g of (±)-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[ethoxy[6-methyl-3-(phenylmethoxy)-2-pyridinyl]-methyl]-1*H*-benzimidazol-2-amine trihydrochloride dihydrate (compound 11).

20 Example B11

A mixture of intermediate (12) (0.0016 mol) and benzenemethanamine (0.0048 mol) in methanol (7ml) was hydrogenated at 40°C under a 5 bar pressure for 8 hours with Pd/C (0.07g) as a catalyst. After uptake of H<sub>2</sub> (1 equivalent), the catalyst was filtered through celite, washed with CH<sub>3</sub>OH and CH<sub>2</sub>Cl<sub>2</sub> and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/CH<sub>3</sub>OH/NH<sub>4</sub>OH 93/7/0.7). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from diethyl ether. The precipitate was filtered off and dried, yielding 0.4g of (±)-*N*-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine (59%) (compound 12).

30 Example B12

A mixture of (±)-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(6-methyl-2-pyridinyl) [2-(phenylmethoxy)ethoxy]methyl]-1*H*-benzimidazol-2-amine (0.003 mol) in methanol (150ml) was stirred at room temperature with Pd/C 10% (0.5g) as a catalyst. After uptake of H<sub>2</sub> (1 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent: CH<sub>2</sub>Cl<sub>2</sub>/(CH<sub>3</sub>OH/NH<sub>3</sub>) 90/10). The pure fractions were collected and the solvent was


evaporated. The residue was suspended in petroleum ether. The precipitate was filtered off and dried, yielding 0.23g of ( $\pm$ )-2-[[[1-(2-aminoethyl)-4-piperidiny]-amino]-1*H*-benzimidazol-2-yl](6-methyl-2-pyridinyl)methoxy]ethanol monohydrate (18%) (compound 13).

### 5 Example B13

A mixture of ( $\pm$ )-1-[4-[[1-(2-ethoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-yl]amino]-1-piperidiny]-3-methyl-2-butanone (0.0032 mol) in  $\text{NH}_3/\text{CH}_3\text{OH}$  (200 ml) was hydrogenated for 3 days at 20°C with Rh/ $\text{Al}_2\text{O}_3$  5% (1 g) as a catalyst in the presence of a thiophene solution (2 ml). After uptake of  $\text{H}_2$  (1  
10 equivalent), the catalyst was filtered off and the filtrate was evaporated. The residue was purified by column chromatography over silica gel (eluent:  $\text{CH}_2\text{Cl}_2/(\text{CH}_3\text{OH}/\text{NH}_3)$  .95/5). The pure fractions were collected and the solvent was evaporated. The residue was crystallized from diisopropyl ether, filtered off and dried, yielding 0.58 g of  
15 ( $\pm$ )-*N*-[1-(2-amino-3-methylbutyl)-4-piperidiny]-1-[(2-ethoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine (compound 14).

Tables 4 to 8 list the compounds of formula (I) which were prepared according to one of the above examples.

### 20 Table 4




Comp. No.	Ex. No.	b	c	a	L	R <sup>b</sup>	R <sup>a</sup>	Physical data
1	B1	NH	CH	CH	H	CH <sub>3</sub>	H	mp. 146°C
2	B2	CH <sub>2</sub>	N	CH	H	H	H	mp. 150°C; ethanedioate(1:2)
6	B5	NH	CH	CH	H	H	4-Cl	mp. 210°C
31	B2	NH	N	CH	H	H	H	
32	B5	NH	CH	CH	H	H	2-Cl	
33	B5	NH	CH	CH	H	H	4-OCH <sub>3</sub>	
34	B5	NH	CH	CH	H	H	3-Cl	
35	B5	NH	CH	CH	H	H	2-CH <sub>3</sub>	

-54-

Comp. No.	Ex. No.	b	c	a	L	R <sup>b</sup>	R <sup>a</sup>	Physical data
36	B5	NH	CH	CH	H	H	3-CH <sub>3</sub>	mp. 145°C
37	B5	NH	CH	CH	H	H	3-OCH <sub>3</sub>	mp. 162°C
38	B5	NH	CH	CH	H	H	3-F	mp. >230°C
39	B5	NH	CH	CH	H	H	2-F	mp. 205°C
40	B5	NH	CH	CH	H	H	4-CH <sub>3</sub>	mp. 207°C
47	B6	NH	CH	N	*	H	3-CH <sub>3</sub>	

\* = -(CH<sub>2</sub>)<sub>2</sub>-NH-C(=O)-O-C(CH<sub>3</sub>)<sub>3</sub>

Table 5



Comp. No.	Ex. No.	L	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	Physical data
3	B3a	H	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	**	HCl (1:3); H <sub>2</sub> O (1:2); 2-propanolate(1:1)
4	B3b	H	Br	-C <sub>2</sub> H <sub>5</sub>	H	
5	B4	H	H	-C <sub>2</sub> H <sub>5</sub>	H	
9	B8	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	Br	-C <sub>2</sub> H <sub>5</sub>	H	
11	B10	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	**	HCl (1:3); H <sub>2</sub> O(1:2)
13	B12	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -OH	H	H <sub>2</sub> O (1:1)
15	B1	H	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
16	B1	H	CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>2</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
17	B1	H	CH <sub>3</sub>	-[(CH <sub>2</sub> ) <sub>2</sub> -O] <sub>2</sub> -CH <sub>3</sub>	H	
18	B1	H	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	(A)
19	B1	H	Br	-C <sub>2</sub> H <sub>5</sub>	H	(A)
20	B1	H	Br	-C <sub>2</sub> H <sub>5</sub>	H	(B)
21	B1	H	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	(B)
22	B1	H	-CH <sub>2</sub> -O-CH <sub>3</sub>	-CH <sub>3</sub>	H	
23	B1	H	Phenyl	-(CH <sub>2</sub> ) <sub>2</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
24	B1	H	-N(CH <sub>3</sub> ) <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
25	B1	H	-C(=O)-NH <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
26	B1	H	-C(=O)-N(CH <sub>3</sub> ) <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	




Comp. No.	Ex. No	L	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	Physical data
27	B1	H	CH <sub>3</sub>	H	H	
28	B1	H	CH <sub>3</sub>	H	H	HCl (1:3);H <sub>2</sub> O (1:1)
29	B1	H	CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>2</sub> -O-CH <sub>2</sub> -phenyl	H	
30	B1	H	CH <sub>3</sub>	-(CH <sub>2</sub> ) <sub>2</sub> -O-CH <sub>3</sub>	H	HCl (1:1)
63	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	
64	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	
65	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	H	-C <sub>2</sub> H <sub>5</sub>	H	HCl (1:4);H <sub>2</sub> O (1:1)
66	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	H	-[(CH <sub>2</sub> ) <sub>2</sub> -O] <sub>2</sub> -CH <sub>3</sub>	H	
78	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	phenyl	-C <sub>2</sub> H <sub>4</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	HCl (1:3);H <sub>2</sub> O(1:1)
79	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	-N(CH <sub>3</sub> ) <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	HCl (1:4);H <sub>2</sub> O(1:3)
80	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	H	H	HCl (1:4);H <sub>2</sub> O(1:1)
81	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>2</sub> -phenyl	H	
82	B9	-(CH <sub>2</sub> ) <sub>2</sub> -NH <sub>2</sub>	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	H	

\*\* = -O-CH<sub>2</sub>-phenyl

(A) indicates the first isolated stereoisomeric form

(B) indicates the second isolated stereoisomeric form

5 Table 6




Comp. No.	Ex. No	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	R <sup>d</sup>	Physical data
7	B6	Br	-C <sub>2</sub> H <sub>5</sub>	H	H	
8	B7	Br	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	
41	B6	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	H	
42	B6	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	H	
43	B6	H	-C <sub>2</sub> H <sub>5</sub>	H	H	
44	B6	CH <sub>3</sub>	-[C <sub>2</sub> H <sub>4</sub> -O] <sub>2</sub> -CH <sub>3</sub>	H	H	
45	B6	phenyl	-C <sub>2</sub> H <sub>4</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	H	
46	B6	-N(CH <sub>3</sub> ) <sub>2</sub>	-C <sub>2</sub> H <sub>5</sub>	H	H	
48	B6	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>2</sub> -phenyl	H	H	

Comp. No.	Ex. No.	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	R <sup>d</sup>	Physical data
49	B6	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	H	H	
50	B6	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	O-CH <sub>2</sub> -phenyl	H	
51	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(S)]
52	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(R)]
53	B7	Br	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(S)]
54	B7	Br	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(R)]
55	B7	Br	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(R)]
56	B7	Br	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(S)]
57	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(S)]
59	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH <sub>3</sub>	[(A),(R)]
60	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH <sub>3</sub>	[(A),(S)]
61	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH <sub>3</sub>	[(B),(S)]
62	B7	CH <sub>3</sub>	-C <sub>2</sub> H <sub>5</sub>	H	-CH <sub>3</sub>	[(B),(R)]

(A) indicates the first isolated stereoisomeric form

(B) indicates the second isolated stereoisomeric form

Table 7




Comp. No.	Ex. No.	R <sup>a</sup>	R <sup>b</sup>	Physical Data
10	B9	Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	mp. 184°C
12	B11	CH <sub>3</sub>	-CH <sub>3</sub>	mp. 114°C
58	B7	CH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(R)]; H <sub>2</sub> O (1:1); mp. 60°C; [α] <sub>20</sub> <sup>D</sup> (5.20mg/1ml in methanol) = -131.15
67	B9	CH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(S)]; H <sub>2</sub> O (1:1); mp. 91°C; [α] <sub>20</sub> <sup>D</sup> (4.50mg/1ml in methanol) = +126.44
68	B9	CH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(R)]; H <sub>2</sub> O (1:1); mp. 60°C; [α] <sub>20</sub> <sup>D</sup> (5.42mg/1ml in methanol) = +62.18
69	B9	Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(S)]; mp. 70°C; [α] <sub>20</sub> <sup>D</sup> (4.78mg/1ml in methanol) = +133.26

Comp. No.	Ex. No.	R <sup>a</sup>	R <sup>b</sup>	Physical Data
70	B9	Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(A),(R)]; mp. 60°C; [α] <sub>20</sub> <sup>D</sup> (5.43mg/1ml in methanol) = +66.85
71	B9	Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(R)]; H <sub>2</sub> O (1:1); mp. 60°C; [α] <sub>20</sub> <sup>D</sup> (5.08mg/1ml in methanol) = -136.02
72	B9	Br	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(S)]; [α] <sub>20</sub> <sup>D</sup> (5.00mg/1ml in methanol) = -58.00
73	B9	CH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	[(B),(S)]; H <sub>2</sub> O (1:1); mp. 60°C; [α] <sub>20</sub> <sup>D</sup> (4.37mg/1ml in methanol) = -60.18
74	B9	CH <sub>3</sub>	-CH <sub>3</sub>	[(A),(R)]; H <sub>2</sub> O (1:1); mp. 70°C; [α] <sub>20</sub> <sup>D</sup> (5.00mg/1ml in methanol) = +73.00
75	B9	CH <sub>3</sub>	-CH <sub>3</sub>	[(A),(S)]; H <sub>2</sub> O (1:1); mp.<50°C; [α] <sub>20</sub> <sup>D</sup> (4.60mg/1ml in methanol) = +126.52
76	B9	CH <sub>3</sub>	-CH <sub>3</sub>	[(B),(S)]; H <sub>2</sub> O (1:1); mp.<50°C; [α] <sub>20</sub> <sup>D</sup> (4.69mg/1ml in methanol) = -57.78
77	B9	CH <sub>3</sub>	-CH <sub>3</sub>	[(B),(R)]; H <sub>2</sub> O (1:2); mp.<50°C; [α] <sub>20</sub> <sup>D</sup> (4.74mg/1ml in methanol) = -127.64
83	B11	CH <sub>3</sub>	-CH(CH <sub>3</sub> ) <sub>2</sub>	mp. 110°C

(A) indicates the first isolated stereoisomeric form

(B) indicates the second isolated stereoisomeric form

Table 8



5

Comp. No.	Ex. No.	b	R <sup>a</sup>	R <sup>b</sup>	R <sup>c</sup>	R <sup>d</sup>	Physical Data
14	B13	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	H	H <sub>2</sub> O; mp. 60°C HCl(1:3)/H <sub>2</sub> O(1:3)
84	B13	CH	CH <sub>3</sub>	-[(CH <sub>2</sub> ) <sub>2</sub> -O] <sub>2</sub> -CH <sub>3</sub>	H	H	
85	B13	CH	phenyl	-C <sub>2</sub> H <sub>4</sub> -O-C <sub>2</sub> H <sub>5</sub>	H	H	
86	B13	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	H	H	
87	B7	CH	Br	CH <sub>2</sub> -CH <sub>3</sub>	H	CH <sub>3</sub>	
88	B13	CH	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	Cl	CH <sub>3</sub>	HCl(1:3)/H <sub>2</sub> O(1:3)
89	B13	N	CH <sub>3</sub>	-C <sub>2</sub> H <sub>4</sub> -O-CH <sub>3</sub>	H	CH <sub>3</sub>	

C. Pharmacological exampleExample C1 : In vitro screening for activity against Respiratory Syncytial Virus.

- The percent protection against cytopathology caused by viruses (antiviral activity or
- 5 IC<sub>50</sub>) achieved by tested compounds and their cytotoxicity (CC<sub>50</sub>) were both calculated from dose-response curves. The selectivity of the antiviral effect is represented by the selectivity index (SI), calculated by dividing the CC<sub>50</sub> (cytotoxic dose for 50% of the cells) by the IC<sub>50</sub> (antiviral activity for 50 % of the cells).
- 10 Automated tetrazolium-based colorimetric assays were used for determination of IC<sub>50</sub> and CC<sub>50</sub> of test compounds. Flat-bottom, 96-well plastic microtiter trays were filled with 180 µl of Eagle's Basal Medium, supplemented with 5 % FCS (0% for FLU) and 20 mM Hepes buffer. Subsequently, stock solutions (7.8 x final test concentration) of compounds were added in 45 µl volumes to a series of triplicate wells so as to allow
- 15 simultaneous evaluation of their effects on virus- and mock-infected cells. Five five-fold dilutions were made directly in the microtiter trays using a robot system. Untreated virus controls, and HeLa cell controls were included in each test. Approximately 100 TCID<sub>50</sub> of Respiratory Syncytial Virus was added to two of the three rows in a volume of 50 µl. The same volume of medium was added to the third row to measure the
- 20 cytotoxicity of the compounds at the same concentrations as those used to measure the antiviral activity. After two hours of incubation, a suspension ( $4 \times 10^5$  cells/ml) of HeLa cells was added to all wells in a volume of 50µl. The cultures were incubated at 37°C in a 5% CO<sub>2</sub> atmosphere. Seven days after infection the cytotoxicity and the antiviral activity was examined spectrophotometrically. To each well of the microtiter tray, 25 µl
- 25 of a solution of MTT (3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide) was added . The trays were further incubated at 37°C for 2 hours, after which the medium was removed from each cup. Solubilization of the formazan crystals was achieved by adding 100 µl 2-propanol. Complete dissolution of the formazan crystals were obtained after the trays have been placed on a plate shaker for 10 min. Finally, the
- 30 absorbances were read in an eight-channel computer-controlled photometer (Multiskan MCC, Flow Laboratories) at two wavelengths (540 and 690 nm). The absorbance measured at 690 nm was automatically subtracted from the absorbance at 540 nm, so as to eliminate the effects of non-specific absorption.
- 35 Particular IC<sub>50</sub>, CC<sub>50</sub> and SI values are listed in Table 9 hereinbelow.


Table 9

-59-

Co. No.	IC <sub>50</sub> (μM)	CC <sub>50</sub> (μM)	SI
87	0.00032	10.12	31623
10	0.0006	37.86	63096
88	0.002	20	10000
67	0.004	63.40	15849
13	0.0126	>100.08	>7943
58	0.0501	79.41	1585
11	0.1259	9.95	79
80	1.2589	>99.45	>79

Claims

1. A compound of formula



a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex or stereochemically isomeric form thereof wherein

-a¹=a²-a³=a⁴- represents a bivalent radical of formula

-CH=CH-CH=CH- (a-1);


-N=CH-CH=CH- (a-2);

-CH=N-CH=CH- (a-3);

-CH=CH-N=CH- (a-4); or


-CH=CH-CH=N- (a-5);

wherein each hydrogen atom in the radicals (a-1), (a-2), (a-3), (a-4) and (a-5) may optionally be replaced by halo, C<sub>1-6</sub>alkyl, nitro, amino, hydroxy, C<sub>1-6</sub>alkyloxy, polyhaloC<sub>1-6</sub>alkyl, carboxyl, aminoC<sub>1-6</sub>alkyl, mono- or di(C<sub>1-4</sub>alkyl)-aminoC<sub>1-6</sub>alkyl, C<sub>1-6</sub>alkyloxycarbonyl, hydroxyC<sub>1-6</sub>alkyl, or a radical of formula



wherein =Z is =O, =CH-C(=O)-NR<sup>5a</sup>R<sup>5b</sup>, =CH<sub>2</sub>, =CH-C<sub>1-6</sub>alkyl, =N-OH or =N-O-C<sub>1-6</sub>alkyl;

20 Q is a radical of formula



wherein Alk is C<sub>1-6</sub>alkanediyl;

Y¹ is a bivalent radical of formula -NR²- or -CH(NR²R⁴)-;

25 X¹ is NR⁴, S, S(=O), S(=O)<sub>2</sub>, O, CH<sub>2</sub>, C(=O), C(=CH<sub>2</sub>), CH(OH), CH(CH<sub>3</sub>), CH(OCH<sub>3</sub>), CH(SCH<sub>3</sub>), CH(NR<sup>5a</sup>R<sup>5b</sup>), CH<sub>2</sub>-NR⁴ or NR⁴-CH<sub>2</sub>;

$X^2$  is a direct bond,  $CH_2$ ,  $C(=O)$ ,  $NR^4$ ,  $C_{1-4}alkyl-NR^4$ ,  $NR^4-C_{1-4}alkyl$ ;

$t$  is 2, 3, 4 or 5;

$u$  is 1, 2, 3, 4 or 5;

$v$  is 2 or 3; and

- 5 whereby each hydrogen atom in Alk and the carbocycles and the heterocycles defined in radicals (b-3), (b-4), (b-5), (b-6), (b-7) and (b-8) may optionally be replaced by  $R^3$ ; with the proviso that when  $R^3$  is hydroxy or  $C_{1-6}alkyloxy$ , then  $R^3$  can not replace a hydrogen atom in the  $\alpha$  position relative to a nitrogen atom;

G is  $C_{1-10}alkanediyl$  substituted with one or more hydroxy,  $C_{1-6}alkyloxy$ ,

- 10  $arylC_{1-6}alkyloxy$ ,  $C_{1-6}alkylthio$ ,  $arylC_{1-6}alkylthio$ ,  $HO(-CH_2-CH_2-O)_n$ ,  $C_{1-6}alkyloxy(-CH_2-CH_2-O)_n$  or  $arylC_{1-6}alkyloxy(-CH_2-CH_2-O)_n$ ;

$R^1$  is a monocyclic heterocycle or aryl; said heterocycle being selected from piperidinyl, piperazinyl, pyridyl, pyrazinyl, pyridazinyl, pyrimidinyl, furanyl, tetrahydrofuranyl, thienyl, pyrrolyl, thiazolyl, oxazolyl, imidazolyl, isothiazolyl, pyrazolyl, isoxazolyl,

- 15 oxadiazolyl; and each heterocycle may optionally be substituted with 1 or where possible more, such as 2, 3 or 4, substituents selected from halo, hydroxy, amino, cyano, carboxy,  $C_{1-6}alkyl$ ,  $C_{1-6}alkyloxy$ ,  $C_{1-6}alkylthio$ ,  $C_{1-6}alkyloxyC_{1-6}alkyl$ , aryl,  $arylC_{1-6}alkyl$ ,  $arylC_{1-6}alkyloxy$ , hydroxy $C_{1-6}alkyl$ , mono-or di( $C_{1-6}alkyl$ )amino, mono-or di( $C_{1-6}alkyl$ )amino $C_{1-6}alkyl$ , polyhalo $C_{1-6}alkyl$ ,  $C_{1-6}alkylcarbonylamino$ ,  
20  $C_{1-6}alkyl-SO_2-NR^{5c}$ ,  $aryl-SO_2-NR^{5c}$ ,  $C_{1-6}alkyloxycarbonyl$ ,  $-C(=O)-NR^{5c}R^{5d}$ ,  $HO(-CH_2-CH_2-O)_n$ , halo( $-CH_2-CH_2-O)_n$ ,  $C_{1-6}alkyloxy(-CH_2-CH_2-O)_n$ ,  $arylC_{1-6}alkyloxy(-CH_2-CH_2-O)_n$  and mono-or di( $C_{1-6}alkyl$ )amino( $-CH_2-CH_2-O)_n$ ; each  $n$  independently is 1, 2, 3 or 4;

$R^2$  is hydrogen, formyl,  $C_{1-6}alkylcarbonyl$ , Hetcarbonyl, pyrrolidinyl, piperidinyl,

- 25 homopiperidinyl,  $C_{3-7}cycloalkyl$  substituted with  $N(R^6)_2$ , or  $C_{1-10}alkyl$  substituted with  $N(R^6)_2$  and optionally with a second, third or fourth substituent selected from amino, hydroxy,  $C_{3-7}cycloalkyl$ ,  $C_{2-5}alkanediyl$ , piperidinyl, mono-or di( $C_{1-6}alkyl$ )amino,  $C_{1-6}alkyloxycarbonylamino$ , aryl and aryloxy;

$R^3$  is hydrogen, hydroxy,  $C_{1-6}alkyl$ ,  $C_{1-6}alkyloxy$ ,  $arylC_{1-6}alkyl$  or  $arylC_{1-6}alkyloxy$ ;

- 30  $R^4$  is hydrogen,  $C_{1-6}alkyl$  or  $arylC_{1-6}alkyl$ ;

$R^{5a}$ ,  $R^{5b}$ ,  $R^{5c}$  and  $R^{5d}$  each independently are hydrogen or  $C_{1-6}alkyl$ ; or

$R^{5a}$  and  $R^{5b}$ , or  $R^{5c}$  and  $R^{5d}$  taken together form a bivalent radical of formula  $-(CH_2)_s-$  wherein  $s$  is 4 or 5;

$R^6$  is hydrogen,  $C_{1-4}alkyl$ , formyl, hydroxy $C_{1-6}alkyl$ ,  $C_{1-6}alkylcarbonyl$  or

- 35  $C_{1-6}alkyloxycarbonyl$ ;

aryl is phenyl or phenyl substituted with 1 or more, such as 2, 3 or 4, substituents selected from halo, hydroxy, C<sub>1-6</sub>alkyl, hydroxyC<sub>1-6</sub>alkyl, polyhaloC<sub>1-6</sub>alkyl, and C<sub>1-6</sub>alkyloxy;

Het is pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl.

5

2. A compound according to claim 1 wherein  $-a^1=a^2=a^3=a^4-$  is a radical of formula (a-1) or (a-2).

10

3. A compound according to claim 1 or 2 wherein R<sup>1</sup> is phenyl optionally substituted with halo, C<sub>1-6</sub>alkyl or C<sub>1-4</sub>alkyloxy; or pyridyl optionally substituted with 1 or more substituents selected from arylC<sub>1-6</sub>alkyloxy, C<sub>1-6</sub>alkyloxyC<sub>1-6</sub>alkyl, aryl, mono-or di(C<sub>1-6</sub>alkyl)amino, C(=O)-NR<sup>5c</sup>R<sup>5d</sup>, halo or C<sub>1-6</sub>alkyl.

15

4. A compound according to any one of claims 1 to 3 wherein G is C<sub>1-4</sub>alkanediyl substituted with hydroxy, C<sub>1-6</sub>alkyloxy, HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, C<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>- or arylC<sub>1-6</sub>alkyloxy(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-.

20

5. A compound according to any one of claims 1 to 4 wherein Q is a radical of formula (b-5) wherein v is 2 and Y<sup>1</sup> is -NR<sup>2</sup>-.

6. A compound according to any one of claims 1 to 5 wherein X<sup>1</sup> is NH or CH<sub>2</sub>.

25

7. A compound according to any one of claims 1 to 6 wherein R<sup>2</sup> is hydrogen or C<sub>1-10</sub>alkyl substituted with NHR<sup>6</sup> wherein R<sup>6</sup> is hydrogen or C<sub>1-6</sub>alkyloxycarbonyl.

30


8. A compound according to claim 1 wherein the compound is [(A),(S)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)-ethoxymethyl]-1H-benzimidazol-2-amine; [(A),(S)]-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine (compound 75); (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-6-chloro-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-4-methyl-1H-benzimidazol-2-amine trihydrochloride trihydrate; [(A),(R)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate; (±)-N-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine; [(A),(S)]-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-2-amine monohydrate; (±)-N-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1H-benzimidazol-

35




- 2-amine; [(A),(R)]-*N*-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine monohydrate; (±)-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-2-benzimidazol-2-amine; (±)-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(2-ethoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine; [(B),(S)] *N*-[1-(2-aminopropyl)-4-piperidinyl]-1-[ethoxy(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine monohydrate; (±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-3-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-7-methyl-3*H*-imidazo[4,5-*b*]pyridin-2-amine; (±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(2-ethoxyethoxy)(6-phenyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine; (±)-*N*-[1-(2-aminoethyl)-4-piperidinyl]-1-[(2-methoxyethoxy)(6-methyl-2-pyridinyl)methyl]-1*H*-benzimidazol-2-amine; (±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-4-methyl-1*H*-benzimidazol-2-amine monohydrate; [(A),(R)]-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-amine; (±)-*N*-[1-(2-amino-3-methylbutyl)-4-piperidinyl]-1-[(6-bromo-2-pyridinyl)ethoxymethyl]-1*H*-benzimidazol-2-amine;
- a prodrug, *N*-oxide, addition salt, quaternary amine, metal complex or stereochemically isomeric form thereof.

9. A compound as claimed in any one of claims 1 to 8 for use as a medicine.
10. A pharmaceutical composition comprising a pharmaceutically acceptable carrier, and as active ingredient a therapeutically effective amount of a compound as described in any one of claims 1 to 8.
11. A process of preparing a composition as claimed in claim 10, characterized in that, a pharmaceutically acceptable carrier is intimately mixed with a therapeutically effective amount of a compound as described in any one of claims 1 to 8.
12. An intermediate of formula




with  $R^1$ , G and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, P being a protective group, and  $Q_1$  being defined as Q according to claim 1 provided that it is devoided of the  $R^2$  or  $R^6$  substituent.

13. An intermediate of formula



with  $R^1$ , G and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $(O=)Q_3$  being a carbonyl derivative of Q, said Q being defined according to claim 1, provided that it is devoided of the  $-NR^2R^4$  or  $-NR^2-$  substituent.


14. An intermediate of formula



with  $R^1$ , Q and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $(O=)G_2$  being a carbonyl derivative of G, said G being defined according to claim 1.

15. A process of preparing a compound as claimed in claim 1, characterized by,


a) reacting an intermediate of formula (II-a) or (II-b) with an intermediate of formula (III)



with  $R^1$ , G, Q and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $W_1$  being a suitable leaving group, in the presence of a suitable base and in a suitable reaction-inert solvent;

b) deprotecting an intermediate of formula (IV)


-65-



with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1,  $H-Q_1$  being defined as Q according to claim 1 provided that  $R^2$  or at least one  $R^6$  substituent is hydrogen, and P being a protective group;

5


- c) deprotecting and reducing an intermediate of formula (IV-a)



with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1,  $H-Q_1$  being defined as Q according to claim 1 provided that  $R^2$  or at least one  $R^6$  substituent is hydrogen,  $Q_{1a}(CH=CH)$  being defined as  $Q_1$  provided that  $Q_1$  comprises an unsaturated bond, and P being a protective group;

10


- d) deprotecting an intermediate of formula (V)



with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $H_2N-Q_2$  being defined as Q according to claim 1 provided that both  $R^6$  substituents are hydrogen or  $R^2$  and  $R^4$  are both hydrogen;


15

- e) deprotecting an intermediate of formula (VI)




with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $H_2N-Q_2$  being defined as Q according to claim 1 provided that both  $R^6$  substituents are hydrogen or  $R^2$  and  $R^4$  are both hydrogen, and P being a protective group;

- 5 f) deprotecting an intermediate of formula (VII) or (VIII)



with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1,  $H-Q_1(OH)$  being defined as Q according to claim 1 provided that  $R^2$  or at least one  $R^6$  substituent is hydrogen and provided that Q comprises a hydroxy moiety,  $H_2N-Q_2(OH)$  being defined as Q according to claim 1 provided that both  $R^6$  substituents are hydrogen or  $R^2$  and  $R^4$  are both hydrogen and provided that Q comprises a hydroxy moiety, and P being a protective group;

- 10 g) amination of an intermediate of formula (IX)




15

with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $H_2N-Q_3H$  being defined as Q according to claim 1 provided that both  $R^6$  substituents are hydrogen or  $R^2$  and  $R^4$  are both hydrogen, and the carbon adjacent to the nitrogen carrying the  $R^6$ , or  $R^2$  and  $R^4$  substituents contains at least one hydrogen, in the presence of a suitable amination reagent;

20


- h) reducing an intermediate of formula (X)

-67-



with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4$  defined as in claim 1, and  $H_2N-CH_2-Q_4$  being defined as Q according to claim 1 provided that Q comprises a  $-CH_2-NH_2$  moiety, in the presence of a suitable reducing agent;


- 5 i) reducing an intermediate of formula (X-a)



with G, and  $-a^1=a^2-a^3=a^4$  defined as in claim 1,  $H_2N-CH_2-Q_4$  being defined as Q according to claim 1 provided that Q comprises a  $-CH_2-NH_2$  moiety, and  $R^1$  being defined as  $R^1$  according to claim 1 provided that it comprises at least one substituent, in the presence of a suitable reducing agent and suitable solvent;

10

- j) amination of an intermediate of formula (XI)




with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4$  defined as in claim 1, and  $H_2N-CH_2-CHOH-CH_2-Q_4$  being defined as Q according to claim 1 provided that Q comprises a

15


$CH_2-CHOH-CH_2-NH_2$  moiety, in the presence of a suitable amination reagent;

- k) reacting an intermediate of formula (XII) with formic acid, formamide and ammonia




with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $H-C(=O)-Q_1$  being defined as Q according to claim 1 provided that  $R^2$  or at least one  $R^6$  substituent is formyl;

- l) amination of an intermediate of formula (XIII) by reaction with an intermediate of formula (XIV)



with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and  $R^{2a}-NH-HQ_5$  being defined as Q according to claim 1 provided that  $R^2$  is other than hydrogen and is represented by  $R^{2a}$ ,  $R^4$  is hydrogen, and the carbon atom adjacent to the nitrogen atom carrying the  $R^2$  and  $R^4$  substituents, carries also at least one hydrogen atom, in the presence of a suitable reducing agent;

- m) reducing an intermediate of formula (XV)




with  $R^1$ , G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and


$(R^6)_2N-[(C_{1-9}alkyl)CH_2OH]-NH-HQ_5$  being defined as Q according to claim 1

- provided that  $R^2$  is other than hydrogen and is represented by  $C_{1-10}alkyl$  substituted with  $N(R^6)_2$  and with hydroxy, and the carbon atom carrying the hydroxy, carries also two hydrogen atoms, and provided that  $R^4$  is hydrogen, and the carbon atom adjacent to the nitrogen atom carrying the  $R^2$  and  $R^4$  substituents, carries also at least one hydrogen atom, with a suitable reducing agent;

- n) deprotecting an intermediate of formula (XVI), (XVI-a) or (XVI-b)




-69-




with G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and H-Q<sub>1</sub> being defined as Q according to claim 1 provided that R<sup>2</sup> or at least one R<sup>6</sup> substituent is hydrogen, and R<sup>1a</sup>-(A-O-H)<sub>w</sub>, R<sup>1a'</sup>-(A-O-H)<sub>2</sub> and R<sup>1a''</sup>-(A-O-H)<sub>3</sub> being defined as R<sup>1</sup> according to claim 1 provided that R<sup>1</sup> is substituted with hydroxy, hydroxyC<sub>1-6</sub>alkyl, or HO(-CH<sub>2</sub>-CH<sub>2</sub>-O)<sub>n</sub>-, with w being an integer from 1 to 4 and P or P<sub>1</sub> being a suitable protecting group, with a suitable acid.

o) amination of an intermediate of formula (XVII)



with R<sup>1</sup>, G,  $-a^1=a^2-a^3=a^4-$ , Alk, X<sup>1</sup>, R<sup>2</sup> and R<sup>4</sup> defined as in claim 1, in the presence of a suitable amination agent;

p) amination of an intermediate of formula (XIX)




with R<sup>1</sup>, G, and  $-a^1=a^2-a^3=a^4-$  defined as in claim 1, and Q<sub>6</sub>N-CH<sub>2</sub>-C<sub>1-3</sub>alkyl-NR<sup>4</sup>

-70-


being defined as Q according to claim 1 provided that in the definition of Q, X<sup>2</sup> is C<sub>2-4</sub>alkyl-NR<sup>4</sup>, in the presence of a suitable amination agent;

q) deprotecting an intermediate of formula (XXI)



5 with R<sup>1</sup>, Q, and -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- defined as in claim 1, and HO-G<sub>1</sub> being defined as G according to claim 1 provided that G is substituted with hydroxy or HO-(CH<sub>2</sub>CH<sub>2</sub>O-)<sub>n</sub>;

r) reducing an intermediate of formula (XXII)



10 with R<sup>1</sup>, Q, and -a<sup>1</sup>=a<sup>2</sup>-a<sup>3</sup>=a<sup>4</sup>- defined as in claim 1, and H-G<sub>2</sub>-OH being defined as G according to claim 1 provided that G is substituted with hydroxy and the carbon atom carrying the hydroxy substituent carries also at least one hydrogen, in the presence of a suitable reducing agent.

15 and, if desired, converting compounds of formula (I) into each other following art-known transformations, and further, if desired, converting the compounds of formula (I), into a therapeutically active non-toxic acid addition salt by treatment with an acid, or into a therapeutically active non-toxic base addition salt by treatment with a base, or conversely, converting the acid addition salt form into the  
20 free base by treatment with alkali, or converting the base addition salt into the free acid by treatment with acid; and, if desired, preparing stereochemically isomeric forms, metal complexes, quaternary amines or N-oxide forms thereof.

16. A product containing (a) a compound as defined in claim 1, and (b) another  
25 antiviral compound, as a combined preparation for simultaneous, separate or sequential use in the treatment or the prevention of viral infections.



17. A pharmaceutical composition comprising a pharmaceutically acceptable carrier and as active ingredients (a) a compound as defined in claim 1, and (b) another antiviral compound.

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number  
WO 01/00612 A3(51) International Patent Classification<sup>7</sup>: C07D 401/12,  
A61K 31/437, 31/4465, 31/4545, A61P 11/00, 31/12,  
C07D 471/04, 401/14 // (C07D 471/04, 235:00, 221:00)Pharmaceutica N.V., Turnhoutseweg 30, B-2340 Beerse  
(BE).

(21) International Application Number: PCT/EP00/05675

(74) Agent: QUAGHEBEUR, Luc; Janssen Pharmaceutica  
N.V., Patent Dept. - 3547, Turnhoutseweg 30, B-2340  
Beerse (BE).

(22) International Filing Date: 20 June 2000 (20.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
99202088.3 28 June 1999 (28.06.1999) EP(71) Applicant (for all designated States except US):  
JANSSEN PHARMACEUTICA N.V. [BE/BE]; c/o  
De Corte, Filip - Ext. 3834, Patent Dept., Turnhoutseweg  
30, B-2340 Beerse (BE).(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,  
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,  
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): JANSSENS, Frans,  
Eduard [BE/BE]; Janssen Pharmaceutica N.V., Turnhout-  
seweg 30, B-2340 Beerse (BE). MEERSMAN, Kathleen,  
Petrus, Marie-José [BE/BE]; Janssen Pharmaceutica  
N.V., Turnhoutseweg 30, B-2340 Beerse (BE). SOM-  
MEN, François, Maria [BE/BE]; Janssen Pharmaceutica  
N.V., Turnhoutseweg 30, B-2340 Beerse (BE). ANDRIES,  
Koenraad, Jozef, Lodewijk, Marcel [BE/BE]; Janssen

Published:

— With international search report.

(88) Date of publication of the international search report:  
29 March 2001For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: RESPIRATORY SYNCYTIAL VIRUS REPLICATION INHIBITORS

(57) Abstract: This invention concerns the compounds of formula (I), prodrugs,  $\langle I \rangle$ -N $\langle I \rangle$ -oxides, addition salts, quaternary amines, metal complexes or stereochemically isomeric forms thereof wherein  $-a^1=a^2=a^3=a^4-$  is a radical of formula  $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ ,  $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{N}-\text{CH}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{N}=\text{CH}-$ ,  $-\text{CH}=\text{CH}-\text{CH}=\text{N}-$  wherein each hydrogen atom may optionally be substituted; Q is a radical of formula (b-1), (b-2), (b-3), (b-4), (b-5), (b-6), (b-7), (b-8), wherein Alk is  $\text{C}_{1-6}$ alkanediyl; Y<sup>1</sup> is a bivalent radical of formula  $-\text{NR}^2-$  or  $-\text{CH}(\text{NR}^2\text{R}^4)-$ ; X<sup>1</sup> is  $\text{NR}^4$ , S, S(=O), S(=O)<sub>2</sub>, O, CH<sub>2</sub>, C(=O), CH(=CH<sub>2</sub>), CH(OH), CH(CH<sub>3</sub>), CH(OCH<sub>3</sub>), CH(SCH<sub>3</sub>), CH(NR<sup>5a</sup>R<sup>5b</sup>), CH<sub>2</sub>-NR<sup>4</sup> or NR<sup>4</sup>-CH<sub>2</sub>; X<sup>2</sup> is a direct bond, CH<sub>2</sub>, C(=O), NR<sup>4</sup>,  $\text{C}_{1-6}$ alkyl-NR<sup>4</sup>, NR<sup>4</sup>- $\text{C}_{1-6}$ alkyl; t is 2 to 5; u is 1 to 5; v is 2 or 3; and whereby each hydrogen in Alk and in (b-3), (b-4), (b-5), (b-6), (b-7) and (b-8), may optionally be replaced by R<sup>3</sup>; provided that when R<sup>3</sup> is hydroxy or  $\text{C}_{1-6}$ alkyloxy, then R<sup>3</sup> can not replace a hydrogen atom in the  $\alpha$  position relative to a nitrogen atom; G is substituted  $\text{C}_{1-10}$ alkanediyl wherein the substituent is attached via an oxygen atom; R<sup>1</sup> is an optionally substituted monocyclic heterocycle or aryl; R<sup>2</sup> is hydrogen, formyl,  $\text{C}_{1-6}$ alkylcarbonyl, Hetcarbonyl, pyrrolidinyl, piperidinyl, homopiperidinyl,  $\text{C}_{3-7}$ cycloalkyl or  $\text{C}_{1-10}$ alkyl substituted with N(R<sup>6</sup>)<sub>2</sub> and optionally with another substituent; R<sup>3</sup> is hydrogen, hydroxy,  $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkyloxy, aryl $\text{C}_{1-6}$ alkyl or aryl $\text{C}_{1-6}$ alkyloxy; R<sup>4</sup> is hydrogen,  $\text{C}_{1-6}$ alkyl or aryl $\text{C}_{1-6}$ alkyl; R<sup>5a</sup>, R<sup>5b</sup>, R<sup>5c</sup> and R<sup>5d</sup> are hydrogen or  $\text{C}_{1-6}$ alkyl; or R<sup>5a</sup> and R<sup>5b</sup>, or R<sup>5c</sup> and R<sup>5d</sup> taken together form a bivalent radical of formula  $-(\text{CH}_2)_s-$  wherein s is 4 or 5; R<sup>6</sup> is hydrogen,  $\text{C}_{1-6}$ alkyl, formyl, hydroxy $\text{C}_{1-6}$ alkyl,  $\text{C}_{1-6}$ alkylcarbonyl or  $\text{C}_{1-6}$ alkyloxycarbonyl; aryl is optionally substituted phenyl; Het is pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl; as respiratory syncytial virus replication inhibitors; their preparation, compositions containing them and their use as a medicine.

WO 01/00612 A3

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 00/05675

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 A61K31/437 A61K31/4465 A61K31/4545 A61P11/00  
A61P31/12 C07D471/04 C07D401/14 //(C07D471/04,235:00,  
221:00)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, CHEM ABS Data

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 92 01697 A (JANSSEN PHARMACEUTICA NV) 6 February 1992 (1992-02-06) page 21, line 9 - line 12 -----	1, 10



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*Z\* document member of the same patent family

Date of the actual completion of the international search

14 December 2000

Date of mailing of the international search report

02/01/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel: (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3018

Authorized officer

Alfaro Faus, I

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/EP 00/05675

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9201697 A	06-02-1992	AT 188477 T	15-01-2000
		AU 646280 B	17-02-1994
		AU 8209391 A	18-02-1992
		CA 2086546 A	20-01-1992
		CN 1058216 A, B	29-01-1992
		CS 9102239 A	19-02-1992
		DE 69131895 D	10-02-2000
		DE 69131895 T	20-07-2000
		DK 539420 T	29-05-2000
		EP 0539420 A	05-05-1993
		ES 2142802 T	01-05-2000
		FI 930199 A	18-01-1993
		GR 3032999 T	31-07-2000
		HU 64066 A	29-11-1993
		IL 98865 A	27-11-1995
		JP 3085707 B	11-09-2000
		KR 206723 B	01-07-1999
		MX 9100307 A	28-02-1992
		NO 304791 B	15-02-1999
		NZ 238863 A	26-03-1993
		PL 170580 B	31-01-1997
		PT 98366 A, B	29-05-1992
		RU 2067978 C	20-10-1996
		SK 280690 B	12-06-2000
		US 5360807 A	01-11-1994
		ZA 9105654 A	31-03-1993